

A

UTILITY PATENT APPLICATION TRANSMITTAL

Only for new and continuation-in-part nonprovisional applications under 37 CFR 1.53(b))

Docket No.: P-IX 2965

Total Pages 2

Check box, if applicable:

☐ DUPLICATE

Address to: ASSISTANT COMMISSIONER FOR PATENTS
Box Patent Application
Washington, D.C. 20231

CERTIFICATE OF MAILING BY "EXPRESS MAIL"

"EXPRESS MAIL" MAILING NUMBER EM508279514USDATE OF DEPOSIT JANUARY 30, 1998

I HEREBY CERTIFY THAT THIS PAPER OR FEE IS BEING DEPOSITED WITH THE UNITED STATES POSTAL SERVICE "EXPRESS MAIL POST OFFICE TO ADDRESSEE" SERVICE UNDER 37 C.F.R. 1.10 ON THE DATE INDICATED ABOVE, AND IS ADDRESSED TO THE ASSISTANT COMMISSIONER FOR PATENTS, WASHINGTON, D.C. 20231

L.B. ANDERSON

(TYPED OR PRINTED NAME OF PERSON MAILING PAPER OR FEE)

L.B. Anderson

(SIGNATURE OF PERSON MAILING PAPER OR FEE)

This is a request for filing a
_____ new utility patent application under 37 CFR 1.53(b)(1).

X continuation-in-part under CFR 1.53(b)(2) of prior application serial no. 08/791,391, filed January 30, 1997.

Title: ANTI- $\alpha_v\beta_3$ RECOMBINANT HUMAN ANTIBODIES, NUCLEIC ACIDS ENCODING SAME AND METHODS OF USE

Inventors: William D. Huse and Scott M. Glaser

Enclosed are:

X return receipt postcard
X 1 page application cover sheet
X 116 pages of specification (includes claims and abstract)
X 13 sheets of drawing(s)
_____ pages of an executed combination declaration and power of attorney

_____ An executed assignment and cover sheet

_____ An executed small entity statement

_____ Also enclosed: _____

_____ This application is based on prior foreign application(s) No.(s) _____, filed in _____ on _____, respectively, and priority is hereby claimed therefrom.

_____ This application is based on, and claims the benefit of, U.S. Provisional Application No. 60/_____, filed _____, and entitled _____
OR (PICK ONE AND DELETE THE OTHER) This application is based on, and claims the benefit of, U.S. Provisional Application No. 60/_____ (yet to be assigned), filed _____, which was converted from U.S. Serial No. _____, and entitled _____.

The filing fee has been calculated as shown below:

	Number Filed		Number Extra		Rate		Fee	
					Small Entity	Other Entity	Small Entity	Other Entity
Total Claims	79 - 20	=	59	x	\$11	\$22	=	\$
Independent Claims	24 - 3	=	21	x	\$41	\$82	=	\$
Multiple Dependent Claims Presented: _____ Yes _____ X No					\$135	\$270		\$
					BASIC FEE		\$395	\$790
					TOTAL FEE		\$	\$

01/30/98

00016051.013098

Inventors: Huse and Glaser
Docket No.: P-IX 2965
Page 2


- _____ A check in the amount of \$ _____ to cover the filing fee is enclosed.
- X The payment of the filing fee is to be deferred until the Declaration is filed. Do not charge our deposit account.
- _____ The Commissioner is hereby authorized to charge fees under 37 CFR 1.16 and 1.17 which may be required or credit any overpayment to Deposit Account No. A duplicate copy of this sheet is enclosed.
- X The Commissioner is hereby authorized to charge to Deposit Account No. 03-0370 any fees under 37 CFR 1.17 which may be required under 37 CFR 1.136(a)(3) for an extension of time in any concurrent or future reply requiring a petition for extension of time. A duplicate copy of this sheet is enclosed.

Address all future communications to:

Cathryn Campbell
CAMPBELL & FLORES LLP
4370 La Jolla Village Drive, Suite 700
San Diego, California 92122
telephone: (619) 535-9001
facsimile: (619) 535-8949

Respectfully submitted,

Date: January 30, 1998



David A. Gay
Registration No. 39,200
CAMPBELL & FLORES LLP
4370 La Jolla Village Drive
Suite 700
San Diego, California 92122

00016061-013098

CERTIFICATE OF MAILING BY "EXPRESS MAIL"

"EXPRESS MAIL" MAILING LABEL NUMBER EM508279514US

JANUARY 30, 1998

DATE OF DEPOSIT

HEREBY CERTIFY THAT THIS PAPER OR FEE IS BEING DEPOSITED
WITH THE UNITED STATES POSTAL SERVICE "EXPRESS MAIL POST
OFFICE TO ADDRESSEE" SERVICE UNDER 37 C.F.R. 1.10 ON THE
DATE INDICATED ABOVE AND IS ADDRESSED TO THE ASSISTANT
COMMISSIONER FOR PATENTS, WASHINGTON, D.C. 20231.

L.B. ANDERSON

(TYPED OR PRINTED NAME OF PERSON MAILING PAPER OR FEE)

(SIGNATURE OF PERSON MAILING PAPER OR FEE)

A P P L I C A T I O N

for

UNITED STATES LETTERS PATENT

on

ANTI- $\alpha_v\beta_3$ RECOMBINANT HUMAN ANTIBODIES, NUCLEIC ACIDS
ENCODING SAME AND METHODS OF USE

by

William D. Huse

and

Scott M. Glaser

Sheets of Drawings: 13

Docket No.: P-IX 2965

Attorneys

CAMPBELL & FLORES LLP

4370 La Jolla Village Drive, Suite 700
San Diego, California 92122

ANTI- $\alpha_v\beta_3$ RECOMBINANT HUMAN ANTIBODIES, NUCLEIC ACIDS
ENCODING SAME AND METHODS OF USE

This application is a continuation-in-part of United States Serial No. 08/791,391, filed January 30, 1997, the entire contents of which is incorporated herein by reference.

BACKGROUND OF THE INVENTION

The present invention relates generally to integrin mediated diseases and, more particularly, to nucleic acids encoding $\alpha_v\beta_3$ -inhibitory monoclonal antibodies and to CDR grafted $\alpha_v\beta_3$ -inhibitory antibodies for the therapeutic treatment of $\alpha_v\beta_3$ -mediated diseases.

Integrins are a class of cell adhesion receptors that mediate both cell-cell and cell-extracellular matrix adhesion events. Integrins consist of heterodimeric polypeptides where a single α chain polypeptide noncovalently associates with a single β chain. There are now about 14 distinct α chain polypeptides and at least about 8 different β chain polypeptides which constitute the integrin family of cell adhesion receptors. In general, different binding specificities and tissue distributions are derived from unique combinations of the α and β chain polypeptides or integrin subunits. The family to which a particular integrin is associated with is usually characterized by the β subunit. However, the ligand binding activity of the integrin is largely influenced by the α subunit. For example, vitronectin binding integrins contain the α_v integrin subunit.

It is now known that the vitronectin binding integrins consist of at least three different α_v

Many integrins mediate their cell adhesive functions by recognizing the tripeptide sequence Arg-Gly-Asp (RGD) found within a large number of extracellular matrix proteins. A variety of approaches have attempted to model agents after this sequence to target a particular integrin-mediated pathology. Such approaches include, for example, the use of RGD-containing peptides and peptide analogues which rely on specificity to be conferred by the sequences flanking the RGD core tripeptide sequence. Although there has been some limited success, most RGD-based inhibitors have been shown to be, at most, selective for the targeted integrin and therefore exhibit some cross-reactivity to other non-targeted integrins. Such cross-reactive inhibitors therefore lack the specificity required for use as an efficacious therapeutic. This is particularly true for previously identified inhibitors of the integrin $\alpha_v\beta_3$.

Monoclonal antibodies on the other hand exhibit the specificity required to be used as an effective therapeutic. Antibodies also have the advantage in that they can be routinely generated against essentially any desired antigen. Moreover, with the development of combinatorial libraries, antibodies can now be produced faster and more efficiently than by previously used methods within the art. The use of combinatorial methodology also allows for the selection of the desired antibody along with the simultaneous isolation of the encoding heavy and light chain nucleic acids. Thus, further modification can be performed to the combinatorial antibody without the incorporation of an additional cloning step.

Regardless of the potential advantages associated with the use of monoclonal antibodies as

therapeutics, these molecules nevertheless have the drawback in that they are almost exclusively derived from non-human mammalian organisms. Therefore, their use as therapeutics is limited by the fact that they will

5 normally elicit a host immune response. Methods for substituting the antigen binding site or complementarity determining regions (CDRs) of the non-human antibody into a human framework have been described. Such methods vary in terms of which amino acid residues should be

10 substituted as the CDR as well as which framework residues should be changed to maintain binding specificity. In this regard, it is understood that proper orientation of the β sheet architecture, correct packing of the heavy and light chain interface and

15 appropriate conformation of the CDRs are all important for preserving antigen specificity and affinity within the grafted antibody. However, all of these methods require knowledge of the nucleotide and amino acid sequence of the non-human antibody and the availability

20 of an appropriately modeled human framework.

Thus, there exists a need for the availability of nucleic acids encoding integrin inhibitory antibodies which can be used as compatible therapeutics in humans. For $\alpha_v\beta_3$ -mediated diseases, the present invention

25 satisfies this need and provides related advantages as well.

SUMMARY OF THE INVENTION

The invention provides a Vitaxin antibody and a LM609 grafted antibody exhibiting selective binding

30 affinity to $\alpha_v\beta_3$. The Vitaxin antibody consists of at least one Vitaxin heavy chain polypeptide and at least one Vitaxin light chain polypeptide or functional

fragments thereof. Also provided are the Vitaxin heavy and light chain polypeptides and functional fragments. The LM609 grafted antibody consists of at least one LM609 CDR grafted heavy chain polypeptide and at least one LM609 CDR grafted light chain polypeptide or functional fragment thereof. The invention additionally provides a high affinity LM609 grafted antibody comprising one or more CDRs having at least one amino acid substitution, where the $\alpha_v\beta_3$ binding activity of the high affinity LM609 grafted antibody is enhanced. Nucleic acids encoding Vitaxin and LM609 grafted heavy and light chains as well as nucleic acids encoding the parental non-human antibody LM609 are additionally provided. Functional fragments of such encoding nucleic acids are similarly provided. The invention also provides a method of inhibiting a function of $\alpha_v\beta_3$. The method consists of contacting $\alpha_v\beta_3$ with Vitaxin or a LM609 grafted antibody or functional fragments thereof under conditions which allow binding to $\alpha_v\beta_3$. Finally, the invention provides for a method of treating an $\alpha_v\beta_3$ -mediated disease. The method consists of administering an effective amount of Vitaxin or a LM609 grafted antibody or functional fragment thereof under conditions which allow binding to $\alpha_v\beta_3$.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the nucleotide and deduced amino acid sequence of the variable region of the antibody Vitaxin. Figure 1A shows the nucleotide and deduced amino acid sequences for the Vitaxin heavy chain variable region (Gln1-Ser117; SEQ ID NOS:1 and 2, respectively) while Figure 1B shows the nucleotide and deduced amino acid sequences for the Vitaxin light chain variable region (Glu1-Lys107; SEQ ID NOS:3 and 4, respectively).

Figure 2 shows the nucleotide and deduced amino acid sequence of the variable region of the monoclonal antibody LM609. Figure 2A shows the nucleotide and deduced amino acid sequence of the LM609 heavy chain variable region (SEQ ID NOS:5 and 6, respectively). The variable region extends from amino acid Glu1 to Ala117. Figure 2B shows the nucleotide and deduced amino acid sequence of the LM609 light chain variable region (SEQ ID NOS:7 and 8, respectively). The variable region of the light chain extends from amino acid Asp1 to Lys107.

Figure 3 shows the competitive inhibition of LM609 IgG binding to the integrin $\alpha_{v}\beta_3$ with recombinant LM609 Fab. Soluble recombinant murine LM609 Fab fragments were prepared from periplasmic fractions of M13 bacteriophage clones muLM609M13 12 and muLM609M13 29. The periplasm samples were serially diluted, mixed with either 1 ng/ml, 5 ng/ml, or 50 ng/ml of LM609 IgG and then incubated in 96 well plates coated with purified $\alpha_v\beta_3$. Plates were washed and bound LM609 IgG detected with goat anti-murine Fc specific antibody conjugated to alkaline phosphatase. Fab produced by clone muLM609M13 12 inhibits both 1 ng/ml and 5 ng/ml LM609 IgG binding at all concentrations of Fab greater than 1:27 dilution.

Figure 4 shows the characterization of Vitaxin binding specificity. Figure 4A shows specific binding of Vitaxin to the integrin $\alpha_v\beta_3$ compared to integrins $\alpha_{Ib}\beta_3$ and $\alpha_v\beta_5$. Figure 4B shows the competitive inhibition of LM609 binding to $\alpha_v\beta_3$ by Vitaxin. Figure 4C shows the competitive inhibition of fibrinogen binding to $\alpha_v\beta_3$ by Vitaxin.

Figure 5 shows the inhibition of $\alpha_v\beta_3$ -mediated cell attachment (5A) and migration (5B) by Vitaxin.

Figure 6 shows the reduction in tumor growth due to Vitaxin mediated inhibition of neovascularization. Figure 6A shows the inhibition of the $\alpha_v\beta_3$ -negative Fg and HEp-3 human tumor fragments grown on chick chorioallantoic membranes (CAMs) following Vitaxin treatment. Figure 6B shows the growth inhibition of Vx2 carcinomas implanted subcutaneously in rabbits at two different Vitaxin doses administered 1 day post implantation. Figure 6C similarly shows Vx2 tumor growth inhibition as in Figure 6B, except that four different Vitaxin doses were administered beginning at 7 days post implantation.

Figure 7 shows the nucleotide and deduced amino acid sequence of the light chain variable region of the LM609 grafted antibody fragment (Glu1-Lys107; SEQ ID NOS:31 and 32, respectively). Position 49 of the light chain variable region can at least be either Arg or Met. The nucleotide and deduced amino acid sequence of the heavy chain variable region of the LM609 grafted antibody fragment is shown in Figure 1A (SEQ ID NOS:1 and 2, respectively).

Figure 8 shows the titration of LM609 grafted antibody variants and LM609 grafted Fab on immobilized $\alpha_v\beta_3$. Bacterial cell lysates containing LM609 grafted antibody (closed circles), LM609 grafted antibody variants with improved affinity isolated from the primary libraries (S102, closed squares; Y100, open squares; and Y101, open triangles) or from combinatorial libraries (closed triangles), or an irrelevant Fab (open circles) were titrated on immobilized $\alpha_v\beta_3$.

Figure 9 shows the construction of combinatorial libraries of enhanced LM609 grafted

antibody variants containing multiple amino acid substitutions.

Figure 10 shows the inhibition of fibrinogen binding to $\alpha_v\beta_3$ by LM609 grafted antibody variants.

- 5 Figure 10A shows inhibition of fibrinogen binding to immobilized $\alpha_v\beta_3$. Figure 1B shows correlation of affinity of antibody variants with inhibition of fibrinogen binding.

- 10 Figure 11 shows the inhibition of M21 human melanoma cell adhesion to fibrinogen by LM609 grafted antibody variants. Cell binding to 10 $\mu\text{g/ml}$ fibrinogen-coated substrate was assessed in the presence of various concentrations of LM609 grafted Fab (closed triangles) or the enhanced LM609 grafted Fabs S102 (open circles), G102
15 (closed circles), or C37 (open triangles).

DETAILED DESCRIPTION OF THE INVENTION

- The invention is directed to nucleic acids encoding the monoclonal antibody (MAb) LM609. This antibody specifically recognizes the integrin $\alpha_v\beta_3$ and
20 inhibits its functional activity. The invention is also directed to nucleic acids encoding and to polypeptides comprising non-murine grafted forms of LM609. These grafted antibodies retain the binding specificity and inhibitory activity of the parent murine antibody LM609.
25 The invention is additionally directed to optimized forms of LM609 grafted antibodies that exhibit increased binding affinity and specificity compared to the non-mouse parental forms of the LM609 grafted antibody.

- 30 In one embodiment, the hybridoma expressing LM609 was used as a source to generate and clone cDNAs

encoding LM609. The heavy and light chain encoding cDNAs were sequenced and their CDR regions were substituted into a human antibody framework to generate the non-murine form of the antibody. The substitution or grafting of the CDRs was performed by codon-based mutagenesis to generate a combinatorial antibody Fab library consisting of members that presented alternative residues at certain positions. Screening of the library resulted in the isolation of Vitaxin. As a grafted antibody containing human framework sequences, it is unlikely that Vitaxin will elicit a host immune response and can therefore be advantageously used for the treatment of $\alpha_v\beta_3$ -mediated diseases.

As used herein, the term "monoclonal antibody LM609" or "LM609" is intended to mean the murine monoclonal antibody specific for the integrin $\alpha_v\beta_3$ which is described by Cheresh, D.A. Proc. Natl. Acad. Sci. USA 84:6471-6475 (1987) and by Cheresh and Spiro J. Biol. Chem. 262:17703-17711 (1987). LM609 was produced against and is reactive with the M21 cell adhesion receptor now known as the integrin $\alpha_v\beta_3$. LM609 inhibits the attachment of M21 cells to $\alpha_v\beta_3$ ligands such as vitronectin, fibrinogen and von Willebrand factor (Cheresh and Spiro, *supra*) and is also an inhibitor of $\alpha_v\beta_3$ -mediated pathologies such as tumor induced angiogenesis (Brooks et al. Cell 79:1157-1164 (1994), granulation tissue development in cutaneous wound (Clark et al., Am. J. Pathology, 148:1407-1421 (1996)) and smooth muscle cell migration such as that occurring during restenosis (Choi et al., J. Vascular Surg., 19:125-134 (1994); Jones et al., Proc. Natl. Acad. Sci. 93:2482-2487 (1996)).

As used herein, the term "Vitaxin" is intended to refer to a non-mouse antibody or functional fragment

thereof having substantially the same heavy and light chain CDR amino acid sequences as found in LM609. The term "Vitaxin" when used in reference to heavy or light chain polypeptides is intended to refer to a non-mouse heavy or light chain or functional fragment thereof having substantially the same heavy or light chain CDR amino acid sequences as found in the heavy or light chain of LM609, respectively. When used in reference to a functional fragment, not all LM609 CDRs need to be represented. Rather, only those CDRs that would normally be present in the antibody portion that corresponds to the functional fragment are intended to be referenced as the LM609 CDR amino acid sequences in the Vitaxin functional fragment. Similarly, the use of the term "Vitaxin" in reference to an encoding nucleic acid is intended to refer to a nucleic acid encoding a non-mouse antibody or functional fragment having substantially the same nucleotide sequence as the heavy and light chain CDR nucleotide sequences and encoding substantially the same CDR amino acid sequences as found in LM609.

As used herein, the term "LM609 grafted antibody" is intended to refer to a non-mouse antibody or functional fragment thereof having substantially the same heavy and light chain CDR amino acid sequences as found in LM609 and absent of the substitution of LM609 amino acid residues outside of the CDRs as defined by Kabat et al., U.S. Dept. of Health and Human Services, "Sequences of Proteins of Immunological Interest" (1983). The term "LM609 grafted antibody" or "LM609 grafted" when used in reference to heavy or light chain polypeptides is intended to refer to a non-mouse heavy or light chain or functional fragment thereof having substantially the same heavy or light chain CDR amino acid sequences as found in the heavy or light chain of LM609, respectively, and also

absent of the substitution of LM609 residues outside of the CDRs as defined by Kabat et al., *supra*. When used in reference to a functional fragment, not all LM609 CDRs need to be represented. Rather, only those CDRs that

5 would normally be present in the antibody portion that corresponds to the functional fragment are intended to be referenced as the LM609 CDR amino acid sequences in the LM609 grafted functional fragment. Similarly, the term "LM609 grafted antibody" or "LM609 grafted" used in

10 reference to an encoding nucleic acid is intended to refer to a nucleic acid encoding a non-mouse antibody or functional fragment being absent of the substitution of LM609 amino acids outside of the CDRs as defined by Kabat et al., *supra* and having substantially the same

15 nucleotide sequence as the heavy and light chain CDR nucleotide sequences and encoding substantially the same CDR amino acid sequences as found in LM609 and as defined by Kabat et al., *supra*.

The term "grafted antibody" or "grafted" when

20 used in reference to heavy or light chain polypeptides or functional fragments thereof is intended to refer to a heavy or light chain or functional fragment thereof having substantially the same heavy or light chain CDR of a donor antibody, respectively, and also absent of the

25 substitution of donor amino acid residues outside of the CDRs as defined by Kabat et al., *supra*. When used in reference to a functional fragment, not all donor CDRs need to be represented. Rather, only those CDRs that would normally be present in the antibody portion that

30 corresponds to the functional fragment are intended to be referenced as the donor CDR amino acid sequences in the functional fragment. Similarly, the term "grafted antibody" or "grafted" when used in reference to an encoding nucleic acid is intended to refer to a nucleic

09016061-013099

acid encoding an antibody or functional fragment, being absent of the substitution of donor amino acids outside of the CDRs as defined by Kabat et al., *supra* and having substantially the same nucleotide sequence as the heavy and light chain CDR nucleotide sequences and encoding substantially the same CDR amino acid sequences as found in the donor antibody and as defined by Kabat et al., *supra*.

The meaning of the above terms are intended to include minor variations and modifications of the antibody so long as its function remains uncompromised. Functional fragments such as Fab, F(ab)₂, Fv, single chain Fv (scFv) and the like are similarly included within the definition of the terms LM609 and Vitaxin. Such functional fragments are well known to those skilled in the art. Accordingly, the use of these terms in describing functional fragments of LM609 or the Vitaxin antibody are intended to correspond to the definitions well known to those skilled in the art. Such terms are described in, for example, Harlow and Lane, Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, New York (1989); Molec. Biology and Biotechnology: A Comprehensive Desk Reference (Myers, R.A. (ed.), New York: VCH Publisher, Inc.); Huston et al., Cell Biophysics, 22:189-224 (1993); Plückthun and Skerra, Meth. Enzymol., 178:497-515 (1989) and in Day, E.D., Advanced Immunochimistry, Second Ed., Wiley-Liss, Inc., New York, NY (1990).

As with the above terms used for describing functional fragments of LM609, Vitaxin and a LM609 grafted antibody, the use of terms which reference other LM609, Vitaxin or LM609 grafted antibody domains, functional fragments, regions, nucleotide and amino acid

09016061-013099

sequences and polypeptides or peptides, is similarly intended to fall within the scope of the meaning of each term as it is known and used within the art. Such terms include, for example, "heavy chain polypeptide" or "heavy chain", "light chain polypeptide" or "light chain", "heavy chain variable region" (V_H) and "light chain variable region" (V_L) as well as the term "complementarity determining region" (CDR).

In the case where there are two or more definitions of a term which is used and/or accepted within the art, the definition of the term as used herein is intended to include all such meanings unless explicitly stated to the contrary. A specific example is the use of the term "CDR" to describe the non-contiguous antigen combining sites found within the variable region of both heavy and light chain polypeptides. This particular region has been described by Kabat et al., *supra*, and by Chothia et al., J. Mol. Biol. 196:901-917 (1987) and by MacCallum et al., J. Mol. Biol. 262:732-745 (1996) where the definitions include overlapping or subsets of amino acid residues when compared against each other. Nevertheless, application of either definition to refer to a CDR of LM609, Vitaxin, LM609 grafted antibodies or variants thereof is intended to be within the scope of the term as defined and used herein. The amino acid residues which encompass the CDRs as defined by each of the above cited references are set forth below in Table 1 as a comparison.

09016061.033099

Table 1: CDR Definitions

	<u>Kabat</u> ¹	<u>Chothia</u> ²	<u>MacCallum</u> ³
V _H CDR1	31-35	26-32	30-35
V _H CDR2	50-65	53-55	47-58
5 V _H CDR3	95-102	96-101	93-101
V _L CDR1	24-34	26-32	30-36
V _L CDR2	50-56	50-52	46-55
V _L CDR3	89-97	91-96	89-96

¹ Residue numbering follows the nomenclature of Kabat et
10 al., *supra*

² Residue numbering follows the nomenclature of Chothia et
al., *supra*

³ Residue numbering follows the nomenclature of MacCallum
et al., *supra*

15 As used herein, the term "substantially" or
"substantially the same" when used in reference to a
nucleotide or amino acid sequence is intended to mean
that the nucleotide or amino acid sequence shows a
considerable degree, amount or extent of sequence
20 identity when compared to a reference sequence. Such
considerable degree, amount or extent of sequence
identity is further considered to be significant and
meaningful and therefore exhibit characteristics which
are definitively recognizable or known. Thus, a
25 nucleotide sequence which is substantially the same
nucleotide sequence as a heavy or light chain of LM609,
Vitaxin, or a LM609 grafted antibody including fragments
thereof, refers to a sequence which exhibits
characteristics that are definitively known or
30 recognizable as encoding or as being the amino acid
sequence of LM609, Vitaxin or a LM609 grafted antibody.
Minor modifications thereof are included so long as they

are recognizable as a LM609, Vitaxin or a LM609 grafted antibody sequence. Similarly, an amino acid sequence which is substantially the same amino acid sequence as a heavy or light chain of Vitaxin, a LM609 grafted antibody
5 or functional fragment thereof, refers to a sequence which exhibits characteristics that are definitively known or recognizable as representing the amino acid sequence of Vitaxin or a LM609 grafted antibody and minor modifications thereof.

10 When determining whether a nucleotide or amino acid sequence is substantially the same as Vitaxin or a LM609 grafted antibody, consideration is given to the number of changes relative to the Vitaxin or LM609 grafted antibody together with whether the function is
15 maintained. For example, a single amino acid change in a 3 amino acid CDR or several changes in a 16 amino acid CDR are considered to be substantially the same if $\alpha_V\beta_3$ binding function is maintained. Thus, a nucleotide or amino acid sequence is substantially the same if it
20 exhibits characteristics that are definitively known or recognizable as representing the nucleotide or amino acid sequence of Vitaxin or a LM609 grafted antibody and minor modifications thereof as long as Vitaxin or LM609 grafted antibody function is maintained.

25 As used herein, the term "fragment" when used in reference to a nucleic acid encoding LM609, Vitaxin or a LM609 grafted antibody is intended to mean a nucleic acid having substantially the same sequence as a portion of a nucleic acid encoding LM609, Vitaxin or a LM609
30 grafted antibody. The nucleic acid fragment is sufficient in length and sequence to selectively hybridize to an LM609, a Vitaxin or a LM609 grafted antibody encoding nucleic acid or a nucleotide sequence

that is complementary to an LM609, Vitaxin or LM609 grafted antibody encoding nucleic acid. Therefore, fragment is intended to include primers for sequencing and polymerase chain reaction (PCR) as well as probes for nucleic acid blot or solution hybridization. The meaning of the term is also intended to include regions of nucleotide sequences that do not directly encode LM609 polypeptides such as the introns, and the untranslated region sequences of the LM609 encoding gene.

As used herein, the term "functional fragment" when used in reference to Vitaxin, to a LM609 grafted antibody or to heavy or light chain polypeptides thereof is intended to refer to a portion of Vitaxin or a LM609 grafted antibody including heavy or light chain polypeptides which still retains some or all or the $\alpha_v\beta_3$ binding activity, $\alpha_v\beta_3$ binding specificity and/or integrin $\alpha_v\beta_3$ -inhibitory activity. Such functional fragments can include, for example, antibody functional fragments such as Fab, F(ab)₂, Fv, single chain Fv (scFv). Other functional fragments can include, for example, heavy or light chain polypeptides, variable region polypeptides or CDR polypeptides or portions thereof so long as such functional fragments retain binding activity, specificity or inhibitory activity. The term is also intended to include polypeptides encompassing, for example, modified forms of naturally occurring amino acids such as D-stereoisomers, non-naturally occurring amino acids, amino acid analogues and mimetics so long as such polypeptides retain functional activity as defined above.

As used herein, the term "enhanced" when used in reference to Vitaxin, a LM609 grafted antibody or a functional fragment thereof is intended to mean that a functional characteristic of the antibody has been

altered or augmented compared to a reference antibody so that the antibody exhibits a desirable property or activity. An antibody exhibiting enhanced activity can exhibit, for example, higher affinity or lower affinity binding, or increased or decreased association or dissociation rates compared to a reference antibody. An antibody exhibiting enhanced activity can also exhibit increased stability such as increased half-life in a particular organism. For example, if higher affinity binding is desired, mutations can be introduced into framework or CDR amino acid residues and the resulting antibody variants screened for higher affinity binding to $\alpha_v\beta_3$ relative to a reference antibody such as the LM609 grafted parent antibody.

The invention provides a nucleic acid encoding a heavy chain polypeptide for Vitaxin or a functional fragment thereof. Also provided is a nucleic acid encoding a light chain polypeptide for Vitaxin or a functional fragment thereof. The nucleic acids consist of substantially the same heavy or light chain variable region nucleotide sequences as those shown in Figure 1A and 1B (SEQ ID NOS:1 and 3, respectively) or a fragment thereof.

Vitaxin, including functional fragments thereof, is a non-mouse antibody which exhibits substantially the same binding activity, binding specificity and inhibitory activity as LM609. The Vitaxin Fv Fragment was produced by functionally replacing CDRs within human heavy and light chain variable region polypeptides with the CDRs derived from LM609. Functional replacement of the CDRs was performed by recombinant methods known to those skilled in the art. Such methods are commonly referred to as CDR grafting and

09016061-013098

are the subject matter of U.S. Patent No. 5,225,539. Such methods can also be found described in "Protein Engineering of Antibody Molecules for Prophylactic and Therapeutic Applications in Man," Clark, M. (ed.),
5 Nottingham, England: Academic Titles (1993).

Briefly, LM609 nucleic acid fragments having substantially the same nucleotide and encoding substantially the same amino acid sequence of each of the heavy and light chain CDRs were synthesized and
10 substituted into each of the respective human chain encoding nucleic acids. To maintain functionality of the newly derived Vitaxin antibody, modifications were performed within the non-CDR framework region. These individual changes were made by generating a population
15 of CDR grafted heavy and light chain variable regions wherein all possible changes outside of the CDRs were represented and then selecting the appropriate antibody by screening the population for binding activity. This screen resulted in the selection of the Vitaxin antibody
20 described herein.

The nucleotide sequences of the Vitaxin heavy and light chain variable regions are shown in Figures 1A and 1B, respectively. These sequences correspond substantially to those that encode the heavy and light
25 chain variable region polypeptides of Vitaxin. These Vitaxin nucleic acids are intended to include both the sense and anti-sense strands of the Vitaxin encoding sequences. Single- and double-stranded nucleic acids are similarly included as well as non-coding portions of the
30 nucleic acid such as introns, 5'- and 3'-untranslated regions and regulatory sequences of the gene for example.

As shown in Figure 1A, the Vitaxin heavy chain variable region polypeptide is encoded by a nucleic acid of about 351 nucleotides in length which begins at the amino terminal Gln1 residue of the variable region through to Ser117. This Vitaxin heavy chain variable region encoding nucleic acid is joined to a human IgG1 constant region to yield a coding region of 1431 nucleotides which encodes a heavy chain polypeptide of 477 total amino acids. Shown in Figure 1B is the Vitaxin light chain variable region polypeptide which is encoded by a nucleic acid of about 321 nucleotides in length beginning at the amino terminal Glu1 residue of the variable region through to Lys107. This Vitaxin light chain variable region nucleic acid is joined to a human kappa construct region to yield a coding region of 642 nucleotides which code for a light chain polypeptide of 214 total amino acids.

Minor modification of these nucleotide sequences are intended to be included as heavy and light chain Vitaxin encoding nucleic acids and their functional fragments. Such minor modifications include, for example, those which do not change the encoded amino acid sequence due to the degeneracy of the genetic code as well as those which result in only a conservative substitution of the encoded amino acid sequence. Conservative substitutions of encoded amino acids include, for example, amino acids which belong within the following groups: (1) non-polar amino acids (Gly, Ala, Val, Leu, and Ile); (2) polar neutral amino acids (Cys, Met, Ser, Thr, Asn, and Gln); (3) polar acidic amino acids (Asp and Glu); (4) polar basic amino acids (Lys, Arg and His); and (5) aromatic amino acids (Phe, Trp, Tyr, and His). Other minor modifications are included within the nucleic acids encoding Vitaxin heavy and light

chain polypeptides so long as the nucleic acid or encoded polypeptides retain some or all of their function as described herein.

Thus, the invention also provides a nucleic acid encoding a Vitaxin heavy chain or functional fragment thereof wherein the nucleic acid encodes substantially the same heavy chain variable region amino acid sequence of Vitaxin as that shown in Figure 1A (SEQ ID NO:2) or a fragment thereof. Similarly, the invention also provides a nucleic acid encoding a Vitaxin light chain or functional fragment thereof wherein the nucleic acid encodes substantially the same light chain variable region amino acid sequence of Vitaxin as that shown in Figure 1B (SEQ ID NO:4) or a fragment thereof.

In addition to conservative substitutions of amino acids, minor modifications of the Vitaxin encoding nucleotide sequences which allow for the functional replacement of amino acids are also intended to be included within the definition of the term. The substitution of functionally equivalent amino acids encoded by the Vitaxin nucleotide sequences is routine and can be accomplished by methods known to those skilled in the art. Briefly, the substitution of functionally equivalent amino acids can be made by identifying the amino acids which are desired to be changed, incorporating the changes into the encoding nucleic acid and then determining the function of the recombinantly expressed and modified Vitaxin polypeptide or polypeptides. Rapid methods for making and screening multiple simultaneous changes are well known within the art and can be used to produce a library of encoding nucleic acids which contain all possible or all desired changes and then expressing and screening the library for

Vitaxin polypeptides which retain function. Such methods include, for example, codon based mutagenesis, random oligonucleotide synthesis and partially degenerate oligonucleotide synthesis.

5 Codon based mutagenesis is the subject matter of U.S. Patent Nos. 5,264,563 and 5,523,388 and is advantageous for the above procedures since it allows for the production of essentially any and all desired frequencies of encoded amino acid residues at any and all
10 particular codon positions within an oligonucleotide. Such desired frequencies include, for example, the truly random incorporation of all twenty amino acids or a specified subset thereof as well as the incorporation of a predetermined bias of one or more particular amino
15 acids so as to incorporate a higher or lower frequency of the biased residues compared to other incorporated amino acid residues. Random oligonucleotide synthesis and partially degenerate oligonucleotide synthesis can similarly be used for producing and screening for
20 functionally equivalent amino acid changes. However, due to the degeneracy of the genetic code, such methods will incorporate redundancies at a desired amino acid position. Random oligonucleotide synthesis is the coupling of all four nucleotides at each nucleotide
25 position within a codon whereas partially degenerate oligonucleotide synthesis is the coupling of equal portions of all four nucleotides at the first two nucleotide positions, for example, and equal portions of two nucleotides at the third position. Both of these
30 latter synthesis methods can be found described in, for example, Cwirla et al., Proc. Natl. Acad. Sci. USA 87:6378-6382, (1990) and Devlin et al., Science 249:404-406, (1990).

09045061 0109

Identification of amino acids to be changed can be accomplished by those skilled in the art using current information available regarding the structure and function of antibodies as well as available and current information encompassing methods for CDR grafting procedures. For example, CDRs can be identified within the donor antibody by any or all of the criteria specified in Kabat et al., *supra*, Chothia et al., *supra*, and/or MacCallum et al., *supra*, and any or all non-identical amino acid residues falling outside of these CDR sequences can be changed to functionally equivalent amino acids. Using the above described methods known within the art, any or all of the non-identical amino acids can be changed either alone or in combination with amino acids at different positions to incorporate the desired number of amino acid substitutions at each of the desired positions. The Vitaxin polypeptides containing the desired substituted amino acids are then produced and screened for retention or augmentation of function compared to the unsubstituted Vitaxin polypeptides. Production of the substituted Vitaxin polypeptides can be accomplished by, for example, recombinant expression using methods known to those skilled in the art. Those Vitaxin polypeptides which exhibit retention or augmentation of function compared to unsubstituted Vitaxin are considered to contain minor modifications of the encoding nucleotide sequence which result in the functional replacement of one or more amino acids.

The functional replacement of amino acids is beneficial when producing grafted antibodies having human framework sequences since it allows for the rapid identification of equivalent amino acid residues without the need for structural information or the laborious

09016061-013098

procedures necessary to assess and identify which amino acid residues should be considered for substitution in order to successfully transfer binding function from the donor. Moreover, it eliminates the actual step-wise procedures to change and test the amino acids identified for substitution. Essentially, using the functional replacement approach described above, all non-identical amino acid residues between the donor and the human framework can be identified and substituted with any or all other possible amino acid residues at each non-identical position to produce a population of substituted polypeptides containing all possible or all desired permutations and combinations. The population of substituted polypeptides can then be screened for those substituted polypeptides which retain function. Using the codon based mutagenesis procedures described above, the generation of a library of substituted amino acid residues and the screening of functionally replaced residues has been used for the rapid production of grafted therapeutic antibodies as well as for the rapid alteration of antibody affinity. Such procedures are exemplified in, for example, Rosok et al., J. Biol. Chem. 271:22611-22618 (1996) and in Glaser et al., J. Immunol. 149:3903-3913 (1992), respectively.

In addition to framework residues, amino acids in one or more CDRs can be functionally replaced to allow identification of a modified LM609 grafted antibody having enhanced activity. Using the methods described above for framework residues, amino acid substitutions can similarly be introduced into one or more CDRs in an LM609 grafted antibody. The modified LM609 grafted antibody can be tested for binding activity to determine whether $\alpha_v\beta_3$ binding activity is maintained. The modified LM609 grafted antibody can be further tested to determine

0904661-013098

if activity has been enhanced. Functional replacement of amino acid residues in one or more CDRs therefore allows the identification of an enhanced LM609 grafted antibody having a desirable property such as enhanced activity.

5 To generate modified LM609 grafted antibodies and select those with enhanced activity, several approaches can be employed in the selection of the number of residues within a CDR to mutate as well as the number of CDRs within a LM609 grafted antibody to modify. The
10 choice of selection criteria for mutagenesis of CDRs will depend on the need and desired application of the enhanced antibody. For example, one or a few amino acid positions within a single CDR can be modified to contain selected amino acids at that position. Alternatively,
15 the targeted amino acid positions can be modified to contain all possible amino acids at that position. The resultant population of modified antibodies can then be screened for enhanced activity.

 The construction of modified LM609 grafted
20 antibody populations can also be made where all amino acids positions within a CDR have been mutated to contain all possible amino acids and where amino acid positions within multiple CDRs have been modified to contain variant amino acid residues. In this way, populations
25 can be constructed which range from 2 to $>10^7$ unique members. The larger the population, the more efficient will be the selection of an enhanced LM609 grafted antibody since there will be a larger number of different antibodies within the population. However, a small
30 population of modified LM609 grafted antibodies can be made and successfully used for the selection of enhanced LM609 grafted antibodies. The size of the population of modified LM609 grafted antibodies will depend on the need

of a particular application and can be determined by one skilled in the art.

The generation of modified LM609 grafted antibodies can be achieved by introducing amino acid substitutions into one or more CDRs of an LM609 grafted antibody. For example, single amino acid substitutions can be systematically introduced into a CDR by changing a given amino acid in the CDR to any or all amino acids. Amino acid substitutions can also be introduced into all amino acid positions in one or more of the CDRs or in all of the CDRs, generating a population of modified LM609 grafted antibody variants. This population of modified LM609 grafted antibody variants having single amino acid substitutions can be screened to identify those variants that maintain $\alpha_v\beta_3$ binding activity. The variants having $\alpha_v\beta_3$ binding activity can be further characterized to identify those variants having enhanced activity. Such a systematic approach to introducing single amino acid substitutions and generating a population of LM609 grafted antibody variants to screen for enhanced LM609 grafted antibodies having high affinity binding to $\alpha_v\beta_3$ is described in Example VI.

In addition to generating a population of modified LM609 grafted antibody variants, a particular CDR or a particular amino acid in a CDR can be selected to introduce one or more amino acid substitutions. For example, sequence homology or a structural model can be used to identify particular amino acid positions to introduce amino acid substitutions. In this example, only one or a few modified LM609 grafted antibody variants are generated and screened for binding activity to $\alpha_v\beta_3$. One of skill in the art will know or can determine whether it is desirable to generate a large

population of modified LM609 grafted antibody variants or to generate a limited number of modified LM609 grafted antibody variants to screen and identify an enhanced LM609 grafted antibody having enhanced activity.

5 In addition to identifying enhanced LM609 grafted antibodies by generating a population of modified LM609 grafted antibodies having single amino acid substitutions in a CDR and screening for enhanced activity, enhanced LM609 grafted antibody variants can
10 also be generated by combining two or more mutations, each known to independently result in enhanced activity, into a single antibody. When there are more than two mutations, an efficient way to identify combinations of mutations which further augment activity is to construct
15 all possible combinations and permutations and then select for those with enhanced activity. For example, two single mutations in one or more CDRs can be combined to generate a new modified LM609 grafted antibody having two CDR mutations and screened to determine if the $\alpha_v\beta_3$
20 binding activity is increased over that of the single mutants. Similarly, three mutations can be combined and the resulting modified LM609 grafted antibody screened for enhanced binding activity. Using such an approach of combining CDR mutations, a new population of modified
25 LM609 grafted antibody variants can be generated by incorporating all combinations of the single CDR mutations resulting in enhanced activity into new modified LM609 grafted antibody variants and screening to obtain an optimized enhanced LM609 grafted antibody.

30 An iterative, step-wise approach to identifying an enhanced LM609 grafted antibody is advantageous in that it allows the identification of an antibody having optimal binding activity without the need to generate and

screen a large number of modified LM609 grafted antibody variants. For example, using the approach described in Examples VI and VII in which single mutants were identified and combined into a new population of LM609 grafted antibody variants, enhanced LM609 grafted antibodies having higher affinity were identified by generating 2592 unique variants. In contrast, complete randomization of a single eight amino acid residue CDR would require $>10^{10}$ unique variants. Therefore, such an iterative approach allows identification of enhanced LM609 grafted antibodies having enhanced activity such as high affinity binding by generating a relatively small number of unique modified LM609 grafted antibody variants and screening and identifying those enhanced LM609 grafted antibody variants exhibiting high affinity binding.

An iterative, step-wise approach to identifying enhanced LM609 variants can also be performed using additional steps. Instead of generating all combinations of single amino acid mutations, the single amino acid mutations can be combined in pairs to generate all combinations of double mutants and screened for activity. Those double mutants having enhanced activity can be combined with any or all single mutants to generate triple mutants that are screened for enhanced binding activity. Each iterative round of generating modified LM609 grafted antibody variants can incorporate additional single mutations, and the resulting modified LM609 grafted antibodies can be screened for enhanced activity. The step-wise generation of LM609 grafted antibody variants can thus be used to identify an optimized LM609 grafted antibody. Additionally, such an iterative approach also allows for the identification of

numerous enhanced antibodies which exhibit a range of different, enhanced binding activities.

An optimized LM609 grafted antibody can also be referred to as an LM609-like grafted antibody or an $\alpha_v\beta_3$ -specific grafted antibody and is recognizable because the antibody or functional fragments thereof retains the functional characteristics of LM609. For example, enhanced LM609 grafted antibody variants, which have a single amino acid substitution and have enhanced activity, can be identified and correlated with a specific amino acid substitution. These amino acid substitutions can be combined to generate a new modified LM609 grafted antibody that is tested for activity. Such a combination of advantageous CDR amino acid substitutions can result in an optimized LM609 grafted antibody with multiple CDRs having at least one amino acid substitution or a single CDR having multiple amino acid substitutions, where the modified LM609 grafted antibody has enhanced activity.

Enhanced LM609 grafted antibodies, particularly those optimized by functional replacement of amino acid residues in the CDRs, have desirable enhanced properties such as increased affinity. For example, an optimized LM609 grafted antibody having increased affinity will have higher affinity than the parent antibody used for introducing functional replacement of amino acids. Higher affinity is determined relative to a reference antibody having a similar structure. For example, if the optimized LM609 grafted antibody is an intact antibody containing two heavy chains and two light chains, then higher affinity is determined relative to the intact parent LM609 grafted antibody. Similarly, if the optimized LM609 grafted antibody is an Fab, then higher

affinity is determined relative to the Fab of the parent LM609 grafted antibody.

Although it is not necessary to proceed through multiple optimization steps to obtain a high affinity LM609 grafted antibody, in general, the increase in affinity can correlate with the number of modifications within and between CDRs as well as with the number of optimization steps. Therefore, LM609 grafted antibodies will exhibit a variety of ranges. For example, LM609 grafted antibodies having enhanced affinity will have up to about 2-fold higher affinity or greater, generally greater than about 2- to 5-fold higher affinity such as greater than about 4- to 5-fold higher affinity or about 5- to 10-fold higher affinity than the reference antibody. Particularly, a LM609 grafted antibody having enhanced affinity will have greater than about 10- to 50-fold higher affinity, greater than about 50-fold higher affinity, or greater than about 100-fold higher affinity than the reference antibody.

As described above, functional replacement of CDR amino acid residues can be used to identify LM609 grafted antibodies exhibiting higher affinity than a parent LM609 grafted antibody. Methods discussed above or below for introducing minor modifications into Vitaxin or LM609 grafted antibody encoding nucleotide sequences can similarly be used to generate a library of modified LM609 grafted antibody variants, including methods such as codon based mutagenesis, random oligonucleotide synthesis and partially degenerate oligonucleotide synthesis. For example, codon based mutagenesis has been used to generate such a library of modified LM609 grafted antibody variants having single amino acid substitutions (see Example VI).

After generating a library of modified LM609 grafted antibody variants, the variants can be expressed and screened for binding activity to $\alpha_v\beta_3$. Methods well known to those skilled in the art related to determining antibody-antigen interactions are used to screen for modified LM609 grafted antibodies exhibiting binding activity to $\alpha_v\beta_3$ (Harlow and Lane, *supra*). For example, an ELISA method has been used to screen a library of modified LM609 grafted antibody variants to identify those variants that maintained $\alpha_v\beta_3$ binding activity (see Example VI). Only those modified LM609 grafted antibodies that maintain $\alpha_v\beta_3$ binding activity are considered for further characterization.

Modified LM609 grafted antibodies having $\alpha_v\beta_3$ binding activity can be further characterized to determine which modified LM609 grafted antibody has enhanced activity. The type of assay used to assess enhanced activity depends on the particular desired characteristic. For example, if altered binding activity is desired, then binding assays that allow determination of binding affinity are used. Such assays include binding assays, competition binding assays and surface plasmon resonance as described in Example VI.

Introduction of single amino acid substitutions into CDRs of LM609 grafted antibodies can be used to generate a library of modified LM609 grafted antibodies and screen for binding activity to $\alpha_v\beta_3$. Those modified LM609 grafted antibodies exhibiting binding activity to $\alpha_v\beta_3$ can then be further characterized to identify enhanced LM609 grafted antibodies exhibiting enhanced activity such as higher binding affinity. For example, using such an approach, a number of enhanced LM609 grafted antibodies having single amino acid substitutions

were generated using the heavy chain variable region shown in Figure 1a (SEQ ID NO:2) and the light chain variable region shown in Figure 7 (SEQ ID NO:32), and LM609 grafted antibodies were identified displaying 2 to 13-fold improved affinity over the parent LM609 grafted antibody (see Example VI).

Following identification of enhanced LM609 grafted antibodies having a single amino acid substitution, the amino acid mutations can be combined to further enhance activity. Methods discussed above for introducing single amino acid substitutions into CDRs can similarly be applied to combine amino acid substitutions. For example, a combinatorial library of amino acid mutations that resulted in enhanced $\alpha_v\beta_3$ binding affinity was generated using degenerate oligonucleotides and two site hybridization mutagenesis as described in Example VII. Enhanced LM609 grafted antibodies containing multiple CDR amino acid substitutions were generated using the heavy chain variable region shown in Figure 1a (SEQ ID NO:2) and the light chain variable region shown in Figure 7 (SEQ ID NO:32), and LM609 grafted antibodies were identified having 20-fold higher affinity to greater than 90-fold higher affinity than the parent LM609 grafted antibody.

In addition to combining CDR amino acid substitutions to generate an enhanced or optimized LM609 grafted antibody, CDR amino acid substitutions can also be combined with framework mutations that contribute desirable properties to a LM609 grafted antibody. Thus, mutations in CDR or framework regions that enhance activity can be combined to further optimize LM609 grafted antibodies.

The invention further provides fragments of Vitaxin heavy and light chain encoding nucleic acids wherein such fragments consist substantially of the same nucleotide or amino acid sequence as the variable region of Vitaxin heavy or light chain polypeptides. The variable region of the Vitaxin heavy chain polypeptide consists essentially of nucleotides 1-351 and of amino acid residues Gln1 to Ser117 of Figure 1A (SEQ ID NOS:1 and 2, respectively). The variable region of the Vitaxin light chain polypeptide consists essentially of nucleotides 1-321 and of amino acid residues Glu1 to Lys107 of Figure 1B (SEQ ID NOS:3 and 4, respectively). The termini of such variable region encoding nucleic acids is not critical so long as the intended purpose and function remains the same.

Fragments additional to the variable region nucleic acid fragments are provided as well. Such fragments include, for example, nucleic acids consisting substantially of the same nucleotide sequence as a CDR of a Vitaxin heavy or light chain polypeptide. Sequences corresponding to the Vitaxin CDRs include, for example, those regions defined by Kabat et al., *supra*, and/or those regions defined by Chothia et al., *supra*, as well as those defined by MacCallum et al., *supra*. The Vitaxin CDR fragments for each of the above definitions correspond to the nucleotides set forth below in Table 2. The nucleotide sequence numbering is taken from the primary sequence shown in Figures 1A and 1B (SEQ ID NOS:1 and 3) and conforms to the definitions previously set forth in Table 1.

Table 2: Vitaxin CDR Nucleotide Residues

	<u>Kabat</u>	<u>Chothia</u>	<u>MacCallum</u>
V _H CDR1	91-105	76-96	88-105
V _H CDR2	148-198	157-168	139-177
5 V _H CDR3	295-318	298-315	289-315
V _L CDR1	70-102	76-96	88-108
V _L CDR2	148-168	148-156	136-165
V _L CDR3	265-291	271-288	265-288

Similarly, the Vitaxin CDR fragments for each
 10 of the above definitions correspond to the amino acid
 residues set forth below in Table 3. The amino acid
 residue number is taken from the primary sequence shown
 in Figures 1A and 1B (SEQ ID NOS:2 and 4) and conforms to
 the definitions previously set forth in Table 1.

15 **Table 3: Vitaxin CDR Amino Acid Residues**

	<u>Kabat</u>	<u>Chothia</u>	<u>MacCallum</u>
V _H CDR1	Ser31-Ser35	Gly26-Tyr32	Ser30-Ser35
V _H CDR2	Lys50-Gly66	Ser53-Gly56	Trp47-Tyr59
V _H CDR3	His99-Tyr106	Asn100-Ala105	Ala97-Ala105
20 V _L CDR1	Gln24-His34	Ser26-His32	Ser30-Tyr36
V _L CDR2	Tyr50-Ser56	Tyr50-Ser52	Leu46-Ile55
V _L CDR3	Gln89-Thr97	Ser91-His96	Gln89-His96

Thus, the invention also provides nucleic acid
 fragments encoding substantially the same amino acid
 25 sequence as a CDR of a Vitaxin heavy or light chain
 polypeptide.

Nucleic acids encoding Vitaxin heavy and light
 chain polypeptides and fragments thereof are useful for a
 variety of diagnostic and therapeutic purposes. For

example, the Vitaxin nucleic acids can be used to produce Vitaxin antibodies and functional fragments thereof having binding specificity and inhibitory activity against the integrin $\alpha_v\beta_3$. The antibody and functional
5 fragments thereof can be used for the diagnosis or therapeutic treatment of $\alpha_v\beta_3$ -mediated disease. Vitaxin and functional fragments thereof can be used, for example, to inhibit binding activity or other functional activities of $\alpha_v\beta_3$ that are necessary for progression of
10 an $\alpha_v\beta_3$ -mediated disease. Other functional activities necessary for progression of $\alpha_v\beta_3$ -mediated disease include, for example, the activation of $\alpha_v\beta_3$, $\alpha_v\beta_3$ -mediated signal transduction and the $\alpha_v\beta_3$ -mediated prevention of apoptosis. Advantageously, however, Vitaxin comprises
15 non-mouse framework amino acid sequences and as such is less antigenic in regard to the induction of a host immune response. The Vitaxin nucleic acids of the inventions can also be used to model functional equivalents of the encoded heavy and light chain
20 polypeptides.

Thus, the invention provides Vitaxin heavy chain and Vitaxin light chain polypeptides or functional fragments thereof. The Vitaxin heavy chain polypeptide exhibits substantially the same amino acid sequence as
25 that shown in Figure 1A (SEQ ID NO:2) or functional fragment thereof whereas the Vitaxin light chain polypeptide exhibits substantially the same amino acid sequence as that shown in Figure 1B (SEQ ID NO:4) or functional fragment thereof. Also provided is a Vitaxin
30 antibody or functional fragment thereof. The antibody is generated from the above heavy and light chain polypeptides or functional fragments thereof and exhibits selective binding affinity to $\alpha_v\beta_3$.

The invention provides a nucleic acid encoding a heavy chain polypeptide for a LM609 grafted antibody. Also provided is a nucleic acid encoding a light chain polypeptide for a LM609 grafted antibody. The nucleic acids consist of substantially the same heavy chain variable region nucleotide sequence as that shown in Figure 1A (SEQ ID NO:1) and substantially the same light chain variable region nucleotide sequence as that shown in Figure 7 (SEQ ID NO:31) or a fragment thereof.

LM609 grafted antibodies, including functional fragments thereof, are non-mouse antibodies which exhibit substantially the same binding activity, binding specificity and inhibitory activity as LM609. The LM609 grafted antibody Fv fragments described herein are produced by functionally replacing the CDRs as defined by Kabat et al. , hereinafter referred to as "Kabat CDRs," within human heavy and light chain variable region polypeptides with the Kabat CDRs derived from LM609. Functional replacement of the Kabat CDRs is performed by the CDR grafting methods previously described and which is the subject matter of U.S. Patent No. 5,225,539, *supra*. Substitution of amino acid residues outside of the Kabat CDRs can additionally be performed to maintain or augment beneficial binding properties so long as such amino acid substitutions do not correspond to a donor amino acid at that particular position. Such substitutions allow for the modulation of binding properties without imparting any mouse sequence characteristics onto the antibody outside of the Kabat CDRs. Although the production of such antibodies is described herein with reference to LM609 grafted antibodies, the substitution of such non-donor amino acids outside of the Kabat CDRs can be utilized for the production of essentially any grafted antibody. The

The nucleotide sequences of the LM609 grafted antibody heavy and light chain variable regions are shown in Figures 1A and 7, respectively. These sequences correspond substantially to those that encode the heavy and light chain variable region polypeptides of a LM609 grafted antibody. These nucleic acids are intended to include both the sense and anti-sense strands of the LM609 grafted antibody encoding sequences. Single- and double-stranded nucleic acids are similarly included as well as non-coding portions of the nucleic acid such as introns, 5'- and 3'-untranslated regions and regulatory sequences of the gene for example.

Thus, the invention also provides a nucleic acid encoding a LM609 grafted antibody heavy chain

wherein the nucleic acid encodes substantially the same heavy chain variable region amino acid sequence as that shown in Figure 1A (SEQ ID NO:2) or fragment thereof. Similarly, the invention also provides a nucleic acid
5 encoding a LM609 grafted antibody light chain wherein the nucleic acid encodes substantially the same light chain variable region amino acid sequence as that shown in Figure 7 (SEQ ID NO:32) or fragment thereof.

In addition to conservative substitutions of
10 amino acids, minor modifications of the LM609 grafted antibody encoding nucleotide sequences which allow for the functional replacement of amino acids are also intended to be included within the definition of the term. Identification of amino acids to be changed can be
15 accomplished by those skilled in the art using current information available regarding the structure and function of antibodies as well as available and current information encompassing methods for CDR grafting procedures. The substitution of functionally equivalent
20 amino acids encoded by the LM609 grafted antibody nucleotide sequences is routine and can be accomplished by methods known to those skilled in the art. As described previously, such methods include, for example, codon based mutagenesis, random oligonucleotide synthesis
25 and partially degenerate oligonucleotide synthesis and are beneficial when producing grafted antibodies since they allow for the rapid identification of equivalent amino acid residues without the need for structural information.

30 The invention further provides fragments of LM609 grafted antibody heavy and light chain encoding nucleic acids wherein such fragments consist substantially of the same nucleotide or amino acid

09046061-013099

sequence as the variable region of a LM609 grafted antibody heavy or light chain polypeptide. As with Vitaxin, the termini of such variable region encoding nucleic acids is not critical so long as the intended
 5 purpose and function remains the same.

Fragments additional to the variable region nucleic acid fragments are provided as well and include, for example, nucleic acids consisting substantially of the same nucleotide sequence as a CDR of a LM609 grafted
 10 antibody heavy or light chain polypeptide. As with Vitaxin, sequences corresponding to the LM609 grafted antibody CDRs include, for example, those regions defined by Kabat et al., *supra*, Chothia et al., *supra*, as well as those defined by MacCallum et al., *supra*. The LM609
 15 grafted antibody CDR regions will be similar to those described previously for Vitaxin. Moreover, such regions are well known and can be determined by those skilled in the art given the LM609 sequences and teachings provided herein. Thus, the invention also provides nucleic acid
 20 fragments encoding substantially the same amino acid sequence as a CDR of a LM609 grafted antibody heavy or light chain polypeptide.

As with Vitaxin, nucleic acids encoding LM609 grafted antibody heavy and light chain polypeptides and
 25 fragments thereof are useful for a variety of diagnostic and therapeutic purposes. For example, LM609 grafted antibody encoding nucleic acids can be used to produce recombinant antibodies and functional fragments thereof having binding specificity and inhibitory activity
 30 against the integrin $\alpha_v\beta_3$. The antibody and functional fragments thereof can be used for the diagnosis or therapeutic treatment of $\alpha_v\beta_3$ -mediated disease. Such diseases and methods of use for anti- $\alpha_v\beta_3$ antibodies have

been described previously in reference to Vitaxin and are equally applicable to the LM609 grafted antibodies described herein.

Thus, the invention provides LM609 grafted
5 antibody heavy chain and Vitaxin light chain polypeptides
or functional fragments thereof. The LM609 grafted
antibody heavy chain polypeptide exhibits substantially
the same amino acid sequence as that shown in Figure 1A
(SEQ ID NO:2) or functional fragment thereof whereas the
10 LM609 grafted antibody light chain polypeptide exhibits
substantially the same amino acid sequence as that shown
in Figure 7 (SEQ ID NO:32). Also provided is a LM609
grafted antibody or functional fragment thereof. The
antibody is generated from the above heavy and light
15 chain polypeptides or functional fragments thereof and
exhibits selective binding affinity to $\alpha_v\beta_3$.

The invention provides an enhanced LM609
grafted antibody exhibiting selective binding affinity to
 $\alpha_v\beta_3$. The enhanced LM609 grafted antibody contains at
20 least one amino acid substitution in one or more CDRs of
a LM609 grafted heavy chain variable region polypeptide
or a LM609 grafted light chain variable region
polypeptide, wherein the $\alpha_v\beta_3$ binding affinity of the
enhanced LM609 grafted antibody is maintained or
25 enhanced.

To identify enhanced LM609 grafted antibodies,
a library of modified LM609 grafted antibodies was
generated as described above and in Example VI.
Initially, LM609 CDRs were identified and selected to
30 introduce single amino acid substitutions. Utilizing the
numbering system of Kabat et al., *supra*, the CDR residues
selected for mutagenesis were V_H CDR1 Gly-Phe-Thr-Phe-Ser-

04016061.013098

Ser-Trp-Pro-Leu-Thr (SEQ ID NO:90) and Gln-Gln-Ser-Gly-Ser-Trp-Pro-Gln-Thr (SEQ ID NO:92).

The nucleotide sequences encoding the CDRs having single amino acid substitutions were V_H CDR1 GGA
 5 ACT ACC TTC AGT AGC TAT GAC ATG TCT (SEQ ID NO:47), GGA
 TTC ACC TGG AGT AGC TAT GAC ATG TCT (SEQ ID NO:49), and
 GGA TTC ACC TTC CTG AGC TAT GAC ATG TCT (SEQ ID NO:51); V_H
 CDR2 TGG GTC GCA AAA GTT AAA AGT GGT GGT GGT (SEQ ID
 NO:53), AGC ACC TAC TAT CCT GAC ACT GTG CAG GGC (SEQ ID
 10 NO:55), and AGC ACC TAC TAT TTA GAC ACT GTG GAG GGC (SEQ
 ID NO:57); V_H CDR3 GCA AGA CAT AAC CAT GGC AGT TTT GCT TAC
 (SEQ ID NO:59), GCA AGA CAT AAC TAC GGC AGT TAT GCT TAC
 (SEQ ID NO:61), GCA AGA CAT AAC TAC GGC AGT TTT GAT TAC
 (SEQ ID NO:63), GCA AGA CAT AAC TAC GGC AGT TTT TAT TAC
 15 (SEQ ID NO:65), GCA AGA CAT AAC TAC GGC AGT TTT GCT TCT
 (SEQ ID NO:67), GCA AGA CAT AAC TAC GGC AGT TTT GCT ACT
 (SEQ ID NO:69), GCA AGA CAT AAC TAC GGC AGT TTT GCT GAT
 (SEQ ID NO:71), GCA AGA CAT AAC TAC GGC AGT TTT GCT GAG
 (SEQ ID NO:73), GCA AGA CAT AAC TAC GGC AGT TTT GCT ATG
 20 (SEQ ID NO:75), GCA AGA CAT AAC TAC GGC AGT TTT GCT GGG
 (SEQ ID NO:77), and GCA AGA CAT AAC TAC GGC AGT TTT GCT
 GCT (SEQ ID NO:79); V_L CDR1 CAG GCC AGC CAA AGT ATT AGC
 AAC TTT CTA CAC TGG TAT (SEQ ID NO:81); V_L CDR2 CTT CTC
 ATC CGT TAT TCT TCC CAG TCC ATC TCT (SEQ ID NO:83); and V_L
 25 CDR3 CAA CAG AGT AAT AGC TGG CCT CAC ACG (SEQ ID NO:85),
 CAA CAG AGT ACT AGC TGG CCT CAC ACG (SEQ ID NO:87), CAA
 CAG AGT GGC AGC TGG CCT CTG ACG (SEQ ID NO:89) and CAA
 CAG AGT GGC AGC TGG CCT CAG ACG (SEQ ID NO:91).

Enhanced LM609 grafted antibodies having CDRs
 30 with single amino acid substitutions and higher affinity
 binding than the parent LM609 grafted antibody can also
 be identified, where the corresponding amino acid
 mutations are combined to generate new modified LM609

grafted antibodies. Identification is performed by screening for $\alpha_v\beta_3$ binding activity. In some combinations, the LM609 grafted antibody will comprise at least one CDR having two or more amino acid

5 substitutions. The invention provides an enhanced LM609 grafted antibody containing at least one of the following CDRs containing multiple amino acid substitutions: a V_H CDR3 selected from the group consisting of Ala-Arg-His-Asn-His-Gly-Ser-Phe-Ala-Ser (SEQ ID NO:94); Ala-Arg-His-

10 Asn-His-Gly-Ser-Phe-Tyr-Ser (SEQ ID NO:96); Ala-Arg-His-Asn-Tyr-Gly-Ser-Phe-Tyr-Glu (SEQ ID NO:98); and Ala-Arg-His-Asn-Tyr-Gly-Ser-Phe-Tyr-Ser (SEQ ID NO:100).

The nucleotide sequences encoding the CDRs having multiple amino acid substitutions were V_H CDR3 GCA

15 AGA CAT AAC CAT GGC AGT TTT GCT TCT (SEQ ID NO:93), GCA AGA CAT AAC CAT GGC AGT TTT TAT TCT (SEQ ID NO:95), GCA AGA CAT AAC TAC GGC AGT TTT TAT GAG (SEQ ID NO:97), and GCA AGA CAT AAC TAC GGC AGT TTT TAT TCT (SEQ ID NO:99).

The invention also provides an enhanced LM609

20 grafted antibody exhibiting selective binding affinity to $\alpha_v\beta_3$, wherein the enhanced LM609 grafted antibody contains at least one amino acid substitution in two or more CDRs of a LM609 grafted heavy chain variable region polypeptide or a LM609 grafted light chain variable

25 region polypeptide.

An enhanced LM609 grafted antibody containing at least one amino acid substitution in two or more CDRs of a LM609 grafted heavy chain variable region polypeptide or a LM609 grafted light chain variable

30 region polypeptide can include an LM609 grafted antibody containing the combination of CDRs selected from the group consisting of: the V_L CDR1 SEQ ID NO:57 and the V_H

CDR3 SEQ ID NO:50; the V_L CDR1 SEQ ID NO:57, the V_H CDR2
 SEQ ID NO:44 and the V_H CDR3 SEQ ID NO:50; the V_L CDR1 SEQ
 ID NO:57, the V_H CDR2 SEQ ID NO:44 and the V_H CDR3 SEQ ID
 NO:52; the V_L CDR1 SEQ ID NO:57, the V_H CDR2 SEQ ID NO:44
 5 and the V_H CDR3 SEQ ID NO:51; the V_L CDR1 SEQ ID NO:57 and
 the V_H CDR3 SEQ ID NO:52; the V_L CDR3 SEQ ID NO:59, the V_H
 CDR2 SEQ ID NO:44 and the V_H CDR3 SEQ ID NO:50; the V_L
 CDR3 SEQ ID NO:61 and V_H CDR3 SEQ ID NO:50; and the V_L
 CDR3 SEQ ID NO:61, the V_H CDR2 SEQ ID NO:44 and V_H CDR3
 10 SEQ ID NO:50.

In addition to enhanced LM609 grafted
 antibodies containing two or more CDRs having single
 amino acid substitutions, the invention also provides
 enhanced LM609 grafted antibodies wherein at least one of
 15 the CDRs has two or more amino acid substitutions.

Enhanced LM609 grafted antibodies having at
 least one CDR with two or more amino acid substitutions
 can include those containing the combination of CDRs
 selected from the group consisting of: the V_L CDR1 SEQ ID
 20 NO:57, the V_H CDR2 SEQ ID NO:44 and the V_H CDR3 SEQ ID
 NO:63; the V_L CDR3 SEQ ID NO:61, the V_H CDR2 SEQ ID NO:44
 and the V_H CDR3 SEQ ID NO:63; the V_L CDR3 SEQ ID NO:61,
 the V_H CDR2 SEQ ID NO:44 and the V_H CDR3 SEQ ID NO:64; the
 V_L CDR3 SEQ ID NO:61 and the V_H CDR3 SEQ ID NO:63; the V_L
 25 CDR3 SEQ ID NO:61 and the V_H CDR3 SEQ ID NO:65; and the V_L
 CDR3 SEQ ID NO:61, the V_H CDR2 SEQ ID NO:44 and the V_H
 CDR3 SEQ ID NO:66.

The invention additionally provides a high
 affinity LM609 grafted antibody exhibiting selective
 30 binding affinity to $\alpha_v\beta_3$. The high affinity LM609 grafted
 antibody contains at least one amino acid substitution in
 one or more CDRs of a LM609 grafted heavy chain variable

region polypeptide or a LM609 grafted light chain variable region polypeptide, wherein the $\alpha_v\beta_3$ binding affinity of the high affinity LM609 grafted antibody is enhanced.

5 High affinity antibodies can include those containing the combination of CDRs selected from the group consisting of: the V_L CDR1 SEQ ID NO:57 and the V_H CDR3 SEQ ID NO:50; the V_L CDR1 SEQ ID NO:57, the V_H CDR2 SEQ ID NO:44 and the V_H CDR3 SEQ ID NO:50; the V_L CDR1 SEQ ID NO:57, the V_H CDR2 SEQ ID NO:44 and the V_H CDR3 SEQ ID NO:52; the V_L CDR1 SEQ ID NO:57, the V_H CDR2 SEQ ID NO:44 and the V_H CDR3 SEQ ID NO:51; the V_L CDR1 SEQ ID NO:57 and the V_H CDR3 SEQ ID NO:52; the V_L CDR3 SEQ ID NO:59, the V_H CDR2 SEQ ID NO:44 and the V_H CDR3 SEQ ID NO:50; the V_L CDR3 SEQ ID NO:61, the V_H CDR2 SEQ ID NO:44 and the V_H CDR3 SEQ ID NO:63; the V_L CDR3 SEQ ID NO:61 and V_H CDR3 SEQ ID NO:50; the V_L CDR3 SEQ ID NO:61, the V_H CDR2 SEQ ID NO:44 and V_H CDR3 SEQ ID NO:50; the V_L CDR1 SEQ ID NO:57, the V_H CDR2 SEQ ID NO:44 and the V_H CDR3 SEQ ID NO:63; the V_L CDR3 SEQ ID NO:61, the V_H CDR2 SEQ ID NO:44 and the V_H CDR3 SEQ ID NO:64; the V_L CDR3 SEQ ID NO:61 and the V_H CDR3 SEQ ID NO:63; the V_L CDR3 SEQ ID NO:61 and the V_H CDR3 SEQ ID NO:65; and the V_L CDR3 SEQ ID NO:61, the V_H CDR2 SEQ ID NO:44 and the V_H CDR3 SEQ ID NO:66.

25 The invention additionally provides a nucleic acid encoding an enhanced LM609 grafted antibody exhibiting selective binding affinity to $\alpha_v\beta_3$. The enhanced LM609 grafted antibody encoded by the nucleic acid contains at least one amino acid substitution in one or more CDRs of a LM609 grafted heavy chain variable region polypeptide or a LM609 grafted light chain variable region polypeptide, wherein the $\alpha_v\beta_3$ binding

affinity of the enhanced LM609 grafted antibody is maintained or enhanced.

The invention further provides a nucleic acid encoding a high affinity LM609 grafted antibody
5 exhibiting selective binding affinity to $\alpha_v\beta_3$. The high affinity LM609 grafted antibody encoded by the nucleic acid contains at least one amino acid substitution in one or more CDRs of a LM609 grafted heavy chain variable
10 region polypeptide or a LM609 grafted light chain variable region polypeptide, wherein the $\alpha_v\beta_3$ binding affinity of the high affinity LM609 grafted antibody is enhanced.

The invention provides a nucleic acid encoding a heavy chain polypeptide for monoclonal antibody LM609
15 or functional fragment thereof. Also provided is a nucleic acid encoding a light chain polypeptide for monoclonal antibody LM609 or a functional fragment thereof. The nucleic acids consist of substantially the same heavy or light chain variable region nucleotide
20 sequences as that shown in Figure 2A and 2B (SEQ ID NOS:5 and 7, respectively) or a fragment thereof.

As described previously, monoclonal antibody LM609 has been shown in the art to have binding activity to the integrin $\alpha_v\beta_3$. Although specificity can in
25 principle be generated towards essentially any target, LM609 is an integrin inhibitory antibody that exhibits substantial specificity and inhibitory activity to a single member within an integrin family. In this case, LM609 exhibits substantial specificity and inhibitory
30 activity to the $\alpha_v\beta_3$ integrin within the β_3 family. The amino acid or nucleotide sequence of monoclonal antibody

LM609 has never been previously isolated and characterized.

The isolation and characterization of LM609 encoding nucleic acids was performed by techniques known to those skilled in the art and which are described further below in the Examples. Briefly, cDNA from hybridoma LM609 was generated and used as the source for which to isolate LM609 encoding nucleic acids. Isolation was performed by first determining the N-terminal amino acid sequence for each of the heavy and light chain polypeptides and then amplifying by PCR the antibody encoding sequences from the cDNA. The 5' primers were reverse translated to correspond to the newly determined N-terminal amino acid sequences whereas the 3' primers corresponded to sequences substantially similar to antibody constant region sequences. Amplification and cloning of the products resulted in the isolation of the nucleic acids encoding heavy and light chains of LM609.

The nucleotide sequences of the LM609 heavy and light chain variable region sequences are shown in Figure 2A and 2B, respectively. These sequences correspond substantially to those that encode the variable region heavy and light chain polypeptides of LM609. As with the Vitaxin nucleic acids, these LM609 nucleic acids are intended to include both sense and anti-sense strands of the LM609 encoding sequences. Single- and double-stranded nucleic acids are also include as well as non-coding portions of the nucleic acid such as introns, 5'- and 3'-untranslated regions and regulatory sequences of the gene for example.

As shown in Figure 2A, the LM609 heavy chain variable region polypeptide is encoded by a nucleic acid

of about 351 nucleotides in length which begins at the amino terminal Glu1 residue of the variable region through to Ala 117. The murine LM609 antibody heavy chain has an IgG2a constant region. Shown in Figure 2B
5 is the LM609 light chain variable region polypeptide which is encoded by a nucleic acid of about 321 nucleotides in length which begins at the amino terminal Asp1 residue of the variable region through to Lys 107. In the functional antibody, LM609 has a kappa light chain
10 constant region.

As with the Vitaxin nucleic acids, minor modifications of these LM609 nucleotide sequences are intended to be included as heavy and light chain LM609 encoding nucleic acids. Such minor modifications are
15 included within the nucleic acids encoding LM609 heavy and light chain polypeptides so long as the nucleic acids or encoded polypeptides retain some or all of their function as described.

Thus, the invention also provides a nucleic
20 acid encoding a LM609 heavy chain or functional fragment wherein the nucleic acid encodes substantially the same variable region amino acid sequence of monoclonal antibody LM609 as that shown in Figure 2A (SEQ ID NO:6) or a fragment thereof. Similarly, the invention also
25 provides a nucleic acid encoding a LM609 light chain or functional fragment wherein the nucleic acid encodes substantially the same variable region amino acid sequence of monoclonal antibody LM609 as that shown in Figure 2B (SEQ ID NO:8) or a fragment thereof.

30 The invention further provides fragments of LM609 heavy and light chain encoding nucleic acids wherein such fragments consist substantially of the same

5 nucleotide or amino acid sequence as the variable region
of LM609 heavy or light chain polypeptides. The variable
region of the LM609 heavy chain polypeptide consists
essentially of nucleotides 1-351 and of amino acid
10 residues Glu1 to Ala117 of Figure 2A (SEQ ID NOS:5 and 6,
respectively). The variable region of the LM609 light
chain polypeptide consists essentially of nucleotides
1-321 and of amino acid residues Asp1 to Lys107 of Figure
2B (SEQ ID NOS:7 and 8, respectively). The termini of
15 such variable region encoding nucleic acids is not
critical so long as the intended purpose and function
remains the same. Such intended purposes and functions
include, for example, use for the production of
recombinant polypeptides or as hybridization probes for
heavy and light chain variable region sequences.

20 Fragments additional to the variable region
nucleic acid fragments are provided as well. Such
fragments include, for example, nucleic acids consisting
substantially of the same nucleotide sequence as a CDR of
a LM609 heavy or light chain polypeptide. Sequences
corresponding to the LM609 CDRs include, for example,
those regions within the variable region which are
defined by Kabat et al., *supra*, and/or those regions
within the variable regions which are defined by Chothia
25 et al., *supra*, as well as those regions defined by
MacCallum et al., *supra*. The LM609 CDR fragments for
each of the above definitions correspond to the
nucleotides set forth below in Table 4. The nucleotide
sequence numbering is taken from the primary sequence
30 shown in Figures 2A and 2B (SEQ ID NOS:5 and 7) and
conforms to the definitions previously set forth in Table
1.

Table 4: LM609 CDR Nucleotide Residues

	<u>Kabat</u>	<u>Chothia</u>	<u>MacCallum</u>
V _H CDR1	91-105	76-96	88-105
V _H CDR2	148-198	157-168	139-177
5 V _H CDR3	295-318	298-315	288-315
V _L CDR1	70-102	76-96	88-108
V _L CDR2	148-168	148-156	136-165
V _L CDR3	265-291	271-288	265-288

Similarly, the LM609 CDR fragments for each of
 10 the above definitions correspond to the amino acid
 residues set forth below in Table 5. The amino acid
 residue numbering is taken from the primary sequence
 shown in Figures 2A and 2B (SEQ ID NOS:6 and 8) and
 conforms to the definitions set forth in Table 1.

15 **Table 5: LM609 CDR Amino Acid Residues**

	<u>Kabat</u>	<u>Chothia</u>	<u>MacCallum</u>
V _H CDR1	Ser31-Ser35	Gly26-Tyr32	Ser30-Ser35
V _H CDR2	Lys50-Gly66	Ser53-Gly56	Trp47-Tyr59
V _H CDR3	His99-Tyr106	Asn100-Ala105	Ala97-Ala105
20 V _L CDR1	Gln24-His34	Ser26-His32	Ser30-Tyr36
V _L CDR2	Tyr50-Ser56	Tyr50-Ser52	Leu46-Ile55
V _L CDR3	Gln89-Thr97	Ser91-His96	Gln89-His96

Nucleic acids encoding LM609 heavy and light
 chain polypeptides and fragments thereof are useful for a
 25 variety of diagnostic and therapeutic purposes. For
 example, the LM609 nucleic acids can be used to produce
 recombinant LM609 antibodies and functional fragments
 thereof having binding specificity and inhibitory
 activity against the integrin $\alpha_v\beta_3$. The antibody and
 30 functional fragments thereof can be used to determine the

presence or absence of $\alpha_v\beta_3$ in a sample to diagnose the susceptibility or occurrence of an $\alpha_v\beta_3$ -mediated disease. Alternatively, the recombinant LM609 antibodies and functional fragments thereof can be used for the therapeutic treatment of $\alpha_v\beta_3$ -mediated diseases or pathological state. As with Vitaxin, recombinant LM609 and functional fragments thereof can be used to inhibit the binding activity or other functional activities of $\alpha_v\beta_3$ that are necessary for progression of the $\alpha_v\beta_3$ -mediated disease or pathological state.

The LM609 nucleic acids of the invention can also be used to model functional equivalents of the encoded heavy and light chain polypeptides. Such functional equivalents can include, for example, synthetic analogues or mimics of the encoded polypeptides or functional fragments thereof. A specific example would include peptide mimetics of the LM609 CDRs that retain some or substantially the same binding or inhibitory activity of LM609. Additionally, the LM609 encoding nucleic acids can be used to engineer and produce nucleic acids which encode modified forms or derivatives of the antibody LM609, its heavy and light chain polypeptides and functional fragments thereof. As described previously, such modified forms or derivatives include, for example, non-mouse antibodies, their corresponding heavy and light chain polypeptides and functional fragments thereof which exhibit substantially the same binding and inhibitory activity as LM609.

The invention also provides a method of treating an $\alpha_v\beta_3$ -mediated disease. The method consists of administering an effective amount of Vitaxin, a LM609 grafted antibody, an enhanced antibody thereof, or a functional fragment thereof under conditions which allow

binding to $\alpha_v\beta_3$. Also provided is a method of inhibiting a function of $\alpha_v\beta_3$. The method consists of contacting $\alpha_v\beta_3$ with Vitaxin, a LM609 grafted antibody or a functional fragment thereof under conditions which allow binding to $\alpha_v\beta_3$.

As described previously, Vitaxin and LM609 grafted antibodies are monoclonal antibodies which exhibit essentially all of the binding characteristics as does its parental CDR-donor antibody LM609. These characteristics include, for example, significant binding specificity and affinity for the integrin $\alpha_v\beta_3$. The Examples below demonstrate these binding properties and further show that the binding of such antibodies to $\alpha_v\beta_3$ inhibits $\alpha_v\beta_3$ ligand binding and function. Thus, Vitaxin and LM609 grafted antibodies are useful for a large variety of diagnostic and therapeutic purposes directed to the inhibition of $\alpha_v\beta_3$ function.

The integrin $\alpha_v\beta_3$ functions in numerous cell adhesion and migration associated events. As such, the dysfunction or dysregulation of this integrin, its function, or of cells expressing this integrin, is associated with a large number of diseases and pathological conditions. The inhibition $\alpha_v\beta_3$ binding or function can therefore be used to treat or reduce the severity of such $\alpha_v\beta_3$ -mediated pathological conditions. Described below are examples of several pathological conditions mediated by $\alpha_v\beta_3$, since the inhibition of at least this integrin reduces the severity of the condition. These examples are intended to be representative and as such are not inclusive of all $\alpha_v\beta_3$ -mediated diseases. For example, there are numerous pathological conditions additional to those discussed below which exhibit the dysregulation of $\alpha_v\beta_3$ binding,

function or the dysregulation of cells expressing this integrin and in which the pathological condition can be reduced, or will be found to be reduced, by inhibiting the binding $\alpha_v\beta_3$. Such pathological conditions which exhibit this criteria, are intended to be included within the definition of the term as used herein.

Angiogenesis, or neovascularization, is the process where new blood vessels form from pre-existing vessels within a tissue. As described further below, this process is mediated by endothelial cells expressing $\alpha_v\beta_3$ and inhibition of at least this integrin, inhibits new vessel growth. There are a variety of pathological conditions that require new blood vessel formation or tissue neovascularization and inhibition of this process inhibits the pathological condition. As such, pathological conditions that require neovascularization for growth or maintenance are considered to be $\alpha_v\beta_3$ -mediated diseases. The extent of treatment, or reduction in severity, of these diseases will therefore depend on the extent of inhibition of neovascularization. These $\alpha_v\beta_3$ -mediated diseases include, for example, inflammatory disorders such as immune and non-immune inflammation, chronic articular rheumatism, psoriasis, disorders associated with inappropriate or inopportune invasion of vessels such as diabetic retinopathy, neovascular glaucoma and capillary proliferation in atherosclerotic plaques as well as cancer disorders. Such cancer disorders can include, for example, solid tumors, tumor metastasis, angiofibromas, retrolental, fibroplasia, hemangiomas, Kaposi's sarcoma and other cancers which require neovascularization to support tumor growth. Additional diseases which are considered angiogenic include psoriasis and rheumatoid arthritis as well as retinal diseases such as macular degeneration.

Diseases other than those requiring new blood vessels which are $\alpha_v\beta_3$ -mediated diseases include, for example, restenosis and osteoporosis.

Treatment of the $\alpha_v\beta_3$ -mediated diseases can be performed by administering an effective amount of Vitaxin, a LM609 grafted antibody, an enhanced antibody thereof, or a functional fragment thereof so as to bind to $\alpha_v\beta_3$ and inhibit its function. Administration can be performed using a variety of methods known in the art. The choice of method will depend on the specific $\alpha_v\beta_3$ -mediated disease and can include, for example, the *in vivo*, *in situ* and *ex vivo* administration of Vitaxin, a LM609 grafted antibody or functional fragment thereof, to cells, tissues, organs, and organisms. Moreover, such antibodies or functional fragments can be administered to an individual exhibiting or at risk of exhibiting an $\alpha_v\beta_3$ -mediated disease. Definite clinical diagnosis of an $\alpha_v\beta_3$ -mediated disease warrants the administration of Vitaxin, a LM609 grafted antibody or a functional fragment thereof. Prophylactic applications are warranted in diseases where the $\alpha_v\beta_3$ -mediated disease mechanisms precede the onset of overt clinical disease. Thus, individuals with familial history of disease and predicted to be at risk by reliable prognostic indicators can be treated prophylactically to interdict $\alpha_v\beta_3$ -mediated mechanisms prior to their onset.

Vitaxin, a LM609 grafted antibody, an enhanced antibody thereof, or functional fragments thereof can be administered in a variety of formulations and pharmaceutically acceptable media for the effective treatment or reduction in the severity of an $\alpha_v\beta_3$ -mediated disease. Such formulations and pharmaceutically acceptable medias are well known to those skilled in the

art. Additionally, Vitaxin, a LM609 grafted antibody or functional fragments thereof can be administered with other compositions which can enhance or supplement the treatment or reduction in severity of an $\alpha_v\beta_3$ -mediated disease. For example, the coadministration of Vitaxin or a LM609 grafted antibody to inhibit tumor-induced neovascularization and a chemotherapeutic drug to directly inhibit tumor growth is one specific case where the administration of other compositions can enhance or supplement the treatment of an $\alpha_v\beta_3$ -mediated disease.

Vitaxin, a LM609 grafted antibody or functional fragments are administered by conventional methods, in dosages which are sufficient to cause the inhibition of $\alpha_v\beta_3$ integrin binding at the sight of the pathology. Inhibition can be measured by a variety of methods known in the art such as *in situ* immunohistochemistry for the prevalence of $\alpha_v\beta_3$ containing cells at the site of the pathology as well as include, for example, the observed reduction in the severity of the symptoms of the $\alpha_v\beta_3$ -mediated disease.

In vivo modes of administration can include intraperitoneal, intravenous and subcutaneous administration of Vitaxin, a LM609 grafted antibody or a functional fragment thereof. Dosages for antibody therapeutics are known or can be routinely determined by those skilled in the art. For example, such dosages are typically administered so as to achieve a plasma concentration from about 0.01 $\mu\text{g/ml}$ to about 100 $\mu\text{g/ml}$, preferably about 1-5 $\mu\text{g/ml}$ and more preferably about 5 $\mu\text{g/ml}$. In terms of amount per body weight, these dosages typically correspond to about 0.1-300 mg/kg , preferably about 0.2-200 mg/kg and more preferably about 0.5-20 mg/kg . Depending on the need, dosages can be

administered once or multiple times over the course of the treatment. Generally, the dosage will vary with the age, condition, sex and extent of the $\alpha_v\beta_3$ -mediated pathology of the subject and should not be so high as to
5 cause adverse side effects. Moreover, dosages can also be modulated by the physician during the course of the treatment to either enhance the treatment or reduce the potential development of side effects. Such procedures are known and routinely performed by those skilled in the
10 art.

The specificity and inhibitory activity of Vitaxin, LM609 grafted antibodies, an enhanced antibody thereof and functional fragments thereof allow for the therapeutic treatment of numerous $\alpha_v\beta_3$ -mediated diseases.
15 Such diseases include, for example, pathological conditions requiring neovascularization such as tumor growth, and psoriasis as well as those directly mediated by $\alpha_v\beta_3$ such as restenosis and osteoporosis. Thus, the invention provides methods as well as Vitaxin and LM609
20 grafted antibody containing compositions for the treatment of such diseases.

Throughout this application various publications are referenced within parentheses. The disclosures of these publications in their entireties are
25 hereby incorporated by reference in this application in order to more fully describe the state of the art to which this invention pertains.

It is understood that modifications which do not substantially affect the activity of the various
30 embodiments of this invention are also included within the definition of the invention provided herein.

09016061-013068

Accordingly, the following examples are intended to illustrate but not limit the present invention.

EXAMPLE I

Isolation and Characterization of LM609

5

Encoding Nucleic Acids

This Example shows the cloning and sequence determination of LM609 encoding nucleic acids.

LM609 is directed against the human vitronectin receptor, integrin $\alpha_v\beta_3$. $\alpha_v\beta_3$ is highly upregulated in
10 melanoma, glioblastoma, and mammary carcinoma and plays a role in the proliferation of M21 melanoma cells both in vitro and in vivo. $\alpha_v\beta_3$ also plays a role in angiogenesis, restenosis and the formation of granulation tissue in cutaneous wounds. LM609 has been shown to
15 inhibit the adhesion of M21 cells to vitronectin as well as prevent proliferation of M21 cells in vitro. Thus, grafting of LM609 could result in a clinically valuable therapeutic agent.

cdNA Synthesis of LM609 Variable Regions: For
20 cdNA synthesis, total RNA was prepared from 10^8 LM609 hybridoma cells using a modification of the method described by Chomczynski and Sacchi (Chomczynski and Sacchi, *Analyt. Biochem.* 162:156 (1987)). LM609 variable (V) region genes were cloned by reverse
25 transcription-polymerase chain reaction (RT-PCR) and cdNA was synthesized using BRL Superscript kit. Briefly, 5 μ g of total cellular RNA, 650 ng oligo dT and H_2O were brought to a total volume of 55 μ l. The sample was heated to 70°C for 10 min and chilled on ice. Reaction
30 buffer was added and the mixture brought to 10 mM DTT and

1 mM dNTPs and heated at 37°C for 2 minutes. 5 μ l (1000 units) reverse transcriptase was added and incubated at 37°C for 1 hour and then chilled on ice.

All oligonucleotides were synthesized by β -cyanoethyl phosphoramidite chemistry on an ABI 394 DNA synthesizer. Oligonucleotides used for PCR amplification and routine site-directed mutagenesis were purified using oligonucleotide purification cartridges (Applied Biosystems, Foster City, CA). Forward PCR primers were designed from N-terminal protein sequence data generated from purified LM609 antibody. The forward PCR primers contained sequences coding for the first six amino acids in each antibody variable chain (protein sequenced at San Diego State University). The sequence of the light chain forward PCR primer (997) was 5'-GCC CAA CCA GCC ATG GCC GAT ATT GTG CTA ACT CAG-3' (SEQ ID NO:19) whereas the light chain reverse PCR primer (734) was 5'-AC AGT TGG TGC AGC ATC AGC-3' (SEQ ID NO:20) used. This reverse primer corresponds to mouse light chain kappa amino acid residues 109-115. The sequence of the heavy chain forward PCR primer (998) was 5'-ACC CCT GTG GCA AAA GCC GAA GTG CAG CTG GTG GAG-3' (SEQ ID NO:21). Heavy chain reverse PCR primer 733: 5'-GA TGG GGG TGT CGT TTT GGC-3' (SEQ ID NO:22). The PCR primers also contain regions of homology with specific sequences within the immunoexpression vector.

V_L and V_H chains were amplified in two separate 50 μ l reaction mixtures containing 2 μ l of the cDNA-RNA heteroduplex, 66.6 mM Tris-HCl pH 8.8, 1.5 mM $MgCl_2$, 0.2 mM of each four dNTPs, 10 mM 2-mercaptoethanol, 0.25 units Taq polymerase (Boehringer-Mannheim, Indianapolis, IN) and 50 pmoles each of primers 997 and 734 and 998 and

733, respectively. The mixtures were overlaid with mineral oil and cycled for two rounds of PCR with each cycle consisting of 30 seconds at 94°C (denature), 30 seconds at 50°C (anneal), and 30 seconds at 72°C (synthesis). This reaction was immediately followed by 30 cycles of PCR consisting of 30 seconds at 94°C (denature), 30 seconds at 55°C (anneal), and 30 seconds at 72°C (synthesis) followed by a final synthesis reaction for 5 minutes at 72°C. The reaction products were pooled, extracted with CHCl₃ and ethanol precipitated.

Amplified products were resuspended in 20 µl TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8.0) and electrophoresed on a 5% polyacrylamide gel. Bands migrating at expected molecular weights of V_H and V_L were excised, chemically eluted from the gel slice, extracted with organic solvents and ethanol precipitated.

Cloning of amplified V_H and V_L genes into M13 phage immunoexpression vector: The amplified V region gene products were sequentially cloned into the phage immunoexpression vector by hybridization mutagenesis (Near, R. *Biotechniques* 12:88 (1992); Yelton et al., *J. Immunol.* 155:1994-2003 (1995)). Introduction of the amplified V_L and V_H sequences by hybridization mutagenesis positions the antibody sequences in frame with the regulatory elements contained in the M13 vector required for efficient Fab expression. One advantage of this technique is that no restriction endonuclease sites need to be incorporated into the V_L or V_H gene sequences for cloning as is done with conventional DNA ligation methods.

To perform the cloning, 400 ng each of the double-stranded amplified products were first

phosphorylated with polynucleotide kinase. 100 ng of the phosphorylated LM609 V_L product was mixed with 250 ng of uridinylated BS11 phage immunoexpression vector, denatured by heating to 90°C and annealed by gradual

5 cooling to room temperature. BS11 is an M13 immunoexpression vector derived from M13 IX and encodes CH₁ of murine IgG1 and murine kappa light chain constant domain (Huse, W.D. In: Antibody Engineering: A Practical Guide, C.A.K. Borrebaeck, ed. W.H. Freeman and Co.,

10 Publishers, New York, pp. 103-120 (1991)). Nucleotide sequences included in the PCR amplification primers anneal to complementary sequences present in the single-stranded BS11 vector. The annealed mixture was fully converted to a double-stranded molecule with T4 DNA

15 polymerase plus dNTPs and ligated with T4 ligase. 1 µl of the mutagenesis reaction was electroporated into *E. coli* strain DH10B, titered onto a lawn of XL-1 *E. coli* and incubated until plaques formed. Plaque lift assays were performed as described using goat anti-murine kappa

20 chain antibody conjugated to alkaline phosphatase (Yelton et al, *supra*; Huse, W.D., *supra*). Fifteen murine light chain positive M13 phage clones were isolated, pooled and used to prepare uridinylated vector to serve as template for hybridization mutagenesis with the PCR amplified

25 LM609 V_H product.

Clones expressing functional murine LM609 Fab were identified by binding to purified $\alpha_v\beta_3$ by ELISA. Briefly, Immulon II ELISA plates were coated overnight with 1 µg/ml (100 ng/well) $\alpha_v\beta_3$ and nonspecific sites

30 blocked for two hours at 27°C. Soluble Fabs were prepared by isolating periplasmic fractions of cultures of *E. coli* strain MK30-3 (Boehringer Mannheim Co.) infected with the Fab expressing M13 phage clones. Periplasm fractions

were mixed with binding buffer 100 mM NaCl, 50 mM Tris pH 7.4, 2mM CaCl₂, 1 mM MgCl₂, 1 mM MnCl₂, 0.02% NaN₃, 1 mg/ml BSA and incubated with immobilized $\alpha_v\beta_3$ for two hours at 27°C. Plates were washed with binding buffer and bound Fab detected with goat anti-murine kappa chain antibody conjugated to alkaline phosphatase. Four $\alpha_v\beta_3$ reactive clones were identified: muLM609M13 12, 29, 31 and 69. MuLM609M13 12 and 29 gave the strongest signals in the ELISA assay. DNA sequence analysis showed that clones muLM609M13 12, 31 and 69 all had identical light chain sequence and confirmed the previously determined N-terminal amino acid sequence of purified LM609 light chain polypeptide. All four clones had identical V_H DNA sequence and also confirmed the previously determined N-terminal amino acid sequence of purified LM609 heavy chain polypeptide.

To further characterize the binding activity of each clone, soluble Fab fractions were prepared from 50 ml cultures of *E. coli* strain MK30-3 infected with clones 12 and 29 and evaluated for binding to $\alpha_v\beta_3$ in a competitive ELISA with LM609 IgG. The results of this ELISA are shown in Figure 3. Clone muLM609M13 12 was found to inhibit LM609 IgG binding (at LM609 IgG concentrations of 1 ng/ml and 5 ng/ml) to $\alpha_v\beta_3$ in a concentration dependent manner at periplasm titers ranging from neat to 1:80. Clone muLM609M13 12 was plaque purified and both the V region heavy and light chain DNA sequences again determined. Complete DNA sequence of the final clone, muLM609M13 12-5, is shown in Figures 2A and 2B.

EXAMPLE II

Construction of Vitaxin: A CDR GraftedLM609 Functional Fragment

One goal of grafting antibodies is to preserve
5 antibody specificity and affinity when substituting
non-human CDRs into a human antibody framework. Another
goal is to minimize the introduction of foreign amino
acid sequences so as to reduce the possible antigenicity
with a human host. This Example describes procedures for
10 accomplishing both of these goals by producing libraries
of grafted antibodies which represent all
possible members which exhibit the highest affinities for
the desired antigen.

The above library was constructed in *E. coli*
15 wherein the possible CDR and framework changes were
incorporated using codon-based mutagenesis (Kristensson
et al., In: Vaccines 95. Cold Spring Harbor Laboratory
Press. Cold Spring Harbor, NY (1995); Rosok et al., *J.*
Biol. Chem. (271:22611-22613 (1996)). Using these
20 procedures, a library was constructed and a functionally
active humanized anti- $\alpha_v\beta_3$ -inhibitory antibody was
identified.

For the construction of one grafted form of
LM609, human framework sequences showing the highest
25 degree of identity to the murine LM609 V region gene
sequences were selected for receiving the LM609 CDRs.
Human heavy chain V region M72 'CL (HHC30Q, HC Subgroup
3, Kabat et al., *supra*) had 88% identity to frameworks 1,
2 and 3 of LM609 heavy chain and human light chain V
30 region LS1 'CL (HKL312, Kappa subgroup 3, Kabat et al.,
supra) had 79% identity to frameworks 1, 2 and 3 of LM609

light chain. Murine LM609 CDR sequences, as defined by Kabat et al., *supra* were grafted onto the human frameworks. Residues predicted to be buried that might affect the structure and therefore the binding properties of the original murine combining site were taken into consideration when designing possible changes (Singer et al., *supra*; Padlan, E.A. *Mol. Immunol.* 28:489-498 (1991)). This analysis of framework residues considered to be important for preserving the specificity and affinity of the combining site revealed only a few differences. For example, in the heavy chain sequence, the predicted buried residues displayed 100% identity. Of particular note is that Arg16 in human heavy chain V region M72 'CL is a relatively uncommon residue among human chains. However, this residue was also found to be present in LM609 V_H and therefore was retained. Similarly, Arg19 in LM609 is a relatively rare residue among murine heavy chains but it is found to occur in M72 'CL and was therefore retained. In the light chain sequences, two nonidentical buried residues were identified between LM609 and LS1 'CL framework regions at positions 49 and 87. These two positions were therefore incorporated into the grafted antibody library as both human and murine alternatives.

Full-length grafted V region genes were synthesized by PCR using long overlapping oligonucleotides. Light chain oligonucleotides containing mixed amino acid residues at positions 49 and 87 were synthesized as described in Glaser et al. (*J. Immunol.* 149:3903-3913 (1992)) and as illustrated in the oligonucleotides represented as V_L oligo3 and V_L oligo4. (SEQ ID NOS:16 and 17, respectively). All long oligonucleotides were gel purified.

Grafted LM609 heavy and light chain V regions were constructed by mixing 5 overlapping oligonucleotides at equimolar concentrations, in the presence of annealing PCR primers. The heavy chain oligonucleotides map to the following nucleotide positions: V_H oligonucleotide 1 (V_H oligo1), nucleotides (nt) 1-84; (SEQ ID NO:9); V_H oligo2, nt 70-153, (SEQ ID NO:10); V_H oligo3, nt 138-225 (SEQ ID NO:11); V_H oligo4, nt 211-291 (SEQ ID NO:12); V_H oligo5, nt 277-351 (SEQ ID NO:13). Similarly, the Vitaxin light chain oligonucleotides map to the following nucleotide positions: V_L oligonucleotide 1 (V_L oligo1), nucleotides (nt) 1-87; (SEQ ID NO:14); V_L oligo2, nt 73-144, (SEQ ID NO:15); V_L oligo3, nt 130-213 (SEQ ID NO:16); V_L oligo4, nt 199-279 (SEQ ID NO:17); V_L oligo5, nt 265-321 (SEQ ID NO:18). The nucleotide sequences of oligonucleotides used to construct grafted LM609 heavy and light chain variable regions are shown in Table 6. Codon positions 49 and 87 in V_L oligo3, and V_L oligo4 represent the randomized codons. The annealing primers contained at least 18 nucleotide residues complementary to vector sequences for efficient annealing of the amplified V region product to the single-stranded vector. The annealed mixture was fully converted to a double-stranded molecule with T4 DNA polymerase plus dNTPs and ligated with T4 ligase.

0903031-0109
B60E70-73097060

Table 6: Oligonucleotides Used to Construct Grafted LM609 Heavy and Light Chain Variable Regions

	CAGGTGCAGC TGGTGGAGTC TGGGGGAGGC GTTGTGCAGC CTGGAAGGTC CCTGAGACTC	
	TCCTGTGCAG CCTCTGGATT CACC	SEQ ID NO: 9
5	AACTTTTGCG ACCCACTCCA GACCCTTGCC CGGAGCCTGG CGAACCCAAG ACATGTCATA	
	GCTACTGAAG GTGAATCCAG AGGC	SEQ ID NO: 10
	TGGGTCGCAA AAGTTAGTAG TGGTGGTGGT AGCACCTACT ATTTAGACAC TGTGCAGGGC	
	CGATTCACCA TCTCCAGAGA CAATAGT	SEQ ID NO: 11
	TGCACAGTAA TACACGGCTG TGTCTCGGC TCTCAGAGAG TTCATTTGCA GGTATAGGGT	
10	GTTCTTACTA TTGTCTCTGG A	SEQ ID NO: 12
	GTGTATTACT GTGCAAGACA TAACTACGGC AGTTTTGCTT ACTGGGGCCA AGGGACTACA	
	GTGACTGTTT CTAGT	SEQ ID NO: 13
	GAGATTGTGC TAACTCAGTC TCCAGCCACC CTGTCTCTCA GCCCAGGAGA AAGGGCGACT	
	CTTTCCTGCC AGGCCAGCCA AAGTATT	SEQ ID NO: 14
15	GATGAGAAGC CTTGGGGCTT GACCAGGCCT TTGTTGATAC CAGTGTAAGT GGTGCTAAT	
	ACTTTGGCTG GC	SEQ ID NO: 15
	CCAAGGCTTC TCATCWASTA TCGTTCCCAG TCCATCTCTG GGATCCCCGC CAGGTTCACT	
	GGCAGTGGAT CAGGGACAGA TTTC	SEQ ID NO: 16
	GCTGCCACTC TGTTGACAGW AATAGACTGC AAAATCTTCA GGCTCCAGAC TGGAGATAGT	
20	GAGGGTGAAA TCTGTCCCTG A	SEQ ID NO: 17
	CAACAGAGTG GCAGCTGGCC TCACACGTTT GGAGGGGGGA CCAAGGTGGA AATTAAG	
		SEQ ID NO: 18

To generate the library, a portion of the mutagenesis reaction (1 μ l) was electroporated into *E. coli* strain DH10B (BRL), titered onto a lawn of XL-1 (Stratagene, Inc.) and incubated until plaques formed. Replica filter lifts were prepared and plaques containing V_H gene sequences were screened either by hybridization with a digoxigenin-labeled oligonucleotide complementary to LM609 heavy chain CDR 2 sequences or reactivity with 7F11-alkaline phosphatase conjugate, a monoclonal antibody raised against the decapeptide sequence Tyr Pro Tyr Asp Val Pro Asp Tyr Ala Ser (SEQ ID NO:28) appended to the carboxy terminus of the vector CH_1 domain (Biosite,

Inc., San Diego, CA). Fifty clones that were double-positive were pooled and used to prepare uridinylated template for hybridization mutagenesis with the amplified grafted LM609 V_L product.

5 The mutagenesis reaction was performed as described above with the V_H oligonucleotides except that the V_L oligonucleotides 1 to 5 were employed (SEQ ID NOS:14 to 18, respectively). The reaction was electroporated into *E. coli* strain DH10B and filter lifts
10 probed with either goat anti-human kappa chain antibody conjugated to alkaline phosphatase or a goat anti-human Fab antibody using an alkaline phosphatase conjugated rabbit anti-goat secondary reagent for detection. Positive clones co-expressing both V_H and V_L gene
15 sequences were selected (160 total) and used to infect *E. coli* strain MK30-3 for preparing soluble Fab fragments.

 The soluble Fab fragments were screened for binding to $\alpha_v\beta_3$ in an ELISA assay. Four clones that were shown from the ELISA to strongly bind $\alpha_v\beta_3$ were identified
20 and further characterized. These clones were termed huLM609M13-34, 54, 55 and 145. All four clones were plaque purified and three independent subclones from each clone was used to prepare Fab fragments for additional binding analysis to $\alpha_v\beta_3$ by ELISA.

25 In this additional ELISA, duplicate plates were coated with $\alpha_v\beta_3$ ligand and incubated with the huLM609 periplasmic samples. In one plate, bound huLM609 Fab was detected with goat anti-human kappa chain antibody conjugated to alkaline phosphatase and in the other plate
30 bound huLM609 Fab was detected with 7F11-alkaline phosphatase conjugate, the monoclonal antibody

recognizing the decapeptide tag. Subclones huLM609M13-34-1, 2 and 3 and huLM609M13-145-1, 2 and 3 all yielded double positive signals indicating that the Fabs contain functional V_H and V_L polypeptides. These results were confirmed in an ELISA assay on M21 cells, a cell line that expresses the integrin $\alpha_v\beta_3$.

DNA sequence analysis of subclones huLM609M13-34-3 and huLM609M13-145-3 revealed mutations introduced into the library by errors due to oligonucleotide synthesis or by errors arising during PCR amplification. These mutations were corrected in clone huLM609M13-34-3 by site-directed mutagenesis. In the light chain sequence the following corrections were made: His36 to Tyr36 and Lys18 to Arg18. In the heavy chain sequence the following corrections were made: Glu1 to Gln1, Asn3 to Gln3, Leu11 to Val11. Additionally, during the construction of LM609 grafted molecules, residue 28 from the heavy chain was considered to be a non-critical framework residue and the human residue (Thr28) was retained. Subsequently, however, it has been determined that residue 28 can be considered part of the CDR. Therefore, residue 28 was converted to the corresponding mouse residue at that position (Ala28) using site directed mutagenesis with the oligonucleotide 5'-GCT ACT GAA GGC GAA TCC AGA G-3' (SEQ ID NO:29). This change was later determined to not provide benefit over the human framework threonine at this site, and the threonine was retained. The final grafted LM609 clone was designated huLM609M13 1135-4 and is termed herein Vitaxin. The DNA sequence of clone Vitaxin is shown in Figures 2A and 2B.

EXAMPLE III

Functional Characterization of Vitaxin

This Example shows the characterization of Vitaxin's binding specificity, affinity and functional
 5 activity in a number of *in vitro* binding and cell adhesion assays.

The binding specificity of Vitaxin for the integrin $\alpha_v\beta_3$ was initially assessed by measuring binding to $\alpha_v\beta_3$ and its crossreactivity to other α_v - or β_3 -
 10 containing integrins. Specifically, binding specificity was assessed by measuring binding to $\alpha_{IIB}\beta_3$, the major integrin expressed on platelets, and to $\alpha_v\beta_5$, an integrin found prevalent on endothelial cells and connective tissue cell types.

15 Briefly, to determine crossreactivity, integrins were coated onto an ELISA plate and a series of antibody dilutions were measured for Vitaxin binding activity against $\alpha_v\beta_3$ and the other integrins. The integrins $\alpha_v\beta_3$ and $\alpha_v\beta_5$ were isolated by affinity
 20 chromatography as described by Cheresh (1987), *supra*, and Cheresh and Spiro (1987), *supra*. $\alpha_{IIB}\beta_3$ was purchased from CalBiochem. Briefly, an LM609 affinity column (Cheresh and Spiro (1987), *supra*) was used to isolate $\alpha_v\beta_3$ from an octylglucoside human placental lysate, whereas an anti- α_v
 25 affinity column was used to isolate $\alpha_v\beta_5$ from the $\alpha_v\beta_3$ -depleted column flow through. Antibody binding activity was assessed by ELISA using a goat anti-human IgG-alkaline phosphatase conjugate. As a control, a purified human IgG₁ antibody was used since Vitaxin
 30 contains a human IgG₁ backbone.

The results of this assay are shown in Figure 4A and reveal that Vitaxin specifically binds to $\alpha_v\beta_3$ with high affinity. There was no detectable binding to the other α_v - or β_3 -containing integrins at antibody concentrations over 1.0 mg/ml.

In a further series of binding studies, the binding affinity and specificity was assessed in a competitive binding assay with the parental LM609 antibody against $\alpha_v\beta_3$. Competitive binding was measured in an ELISA assay as described above with LM609 being the labeled antibody. Binding of LM609 was determined in the presence of increasing concentrations of Vitaxin competitor. Alternatively, the control competitor antibody was again a human IgG₁.

The results of this competition are presented in Figure 4B and show that specific inhibition of LM609 binding can be observed at Vitaxin concentrations of over 0.1 μ g/ml. Almost complete inhibition is observed at Vitaxin concentrations greater than 100 μ g/ml. This level of competitive inhibition indicates that the parental monoclonal antibody LM609 and the grafted version Vitaxin exhibit essentially identical specificity.

Binding affinity and specificity were also assessed by measuring the inhibitory activity of Vitaxin on $\alpha_v\beta_3$ binding to fibrinogen. For these studies, $\alpha_v\beta_3$ was plated onto ELISA plates as described above for the Vitaxin/ $\alpha_v\beta_3$ binding studies. Inhibitory activity of Vitaxin was determined by measuring the amount of bound biotinylated fibrinogen in the presence of increasing concentrations of Vitaxin or control antibody. Briefly, fibrinogen was purchased from CalBiochem and biotinylated

with N-hydroxysuccinimidobiotin as described by the manufacturer (Pierce Life Science and Analytical Research). Streptavidin alkaline phosphatase was used to detect the bound fibrinogen.

5 The results of this assay are presented in Figure 4C and reveal a specific binding inhibition at Vitaxin concentrations higher than about 0.1 µg/ml. These results, combined with those presented above showing specific binding of Vitaxin to $\alpha_v\beta_3$ and
10 competitive inhibition of LM609, demonstrate that Vitaxin maintains essentially all of the binding characteristics and specificity exhibited by the parental murine monoclonal antibody LM609. Described below are additional functional studies which corroborate these
15 conclusions based on *in vitro* binding assays.

Additional functional studies were performed to further assess the specificity of Vitaxin binding. These studies were directed to the inhibition of integrin $\alpha_v\beta_3$ binding in cell adhesion assays. Endothelial cell
20 adhesion events are an important component in the angiogenic process and inhibition of $\alpha_v\beta_3$ is known to reduce the neovascularization of tumors and thereby reduce the rate of tumor growth. The inhibition of $\alpha_v\beta_3$ -mediated cell attachment by Vitaxin in these assays
25 is indicative of the inhibitory activity expected when this antibody is used *in situ* or *in vivo*.

Briefly, $\alpha_v\beta_3$ -positive M21 melanoma cells grown in RPMI containing 10% FBS were used for these cell binding assays. Cells were released from the culture
30 dish by trypsinization and re-suspended in adhesion buffer at a concentration of 4×10^5 cells/ml (see below). Vitaxin, LM609 or purified human IgG₁ (control antibody),

were diluted to the desired concentration in 250 μ l adhesion buffer (10 mM Hepes, 2 mM $MgCl_2$, 2 mM $CaCl_2$, 0.2 mM $MnCl_2$, and 1% BSA in Hepes buffered saline at pH 7.4) and added to wells of a 48-well plate precoated with
 5 fibrinogen. The fibrinogen was isolated as described above. Each well was coated with 200 μ l fibrinogen at a concentration of 10 μ g/ml for 1 hour at 37°C. For the assay, an equal volume of cells (250 μ l) containing Vitaxin, LM609 or isotype matched control antibody was
 10 added to each of the wells, mixed by gentle shaking and incubated for 20 minutes at 37°C. Unbound cells were removed by washing with adhesion buffer until no cells remained in control wells coated with BSA alone. Bound cells were visualized by staining with crystal violet
 15 which was subsequently extracted with 100 μ l acetic acid (10%) and quantitated by determining the absorbance of the solubilized dye at 560 nm.

The results of this assay are shown in Figure 5A and reveal that both Vitaxin and parental antibody
 20 LM609 inhibit M21 cell adhesion to fibrinogen over the same concentration range. The inhibitory concentration for 50% maximal adhesion was calculated to be about 50 ng/ml. Specificity of Vitaxin was shown by the lack of inhibition observed by the control IgG₁ antibody.

25 In addition to the above cell adhesion results, the inhibitory activity of Vitaxin was also tested in an endothelial cell migration assay. In this regard, the transwell cell migration assay was used to assess the ability of Vitaxin to inhibit endothelial cell migration
 30 (Choi et al., J. Vascular Surg., 19:125-134 (1994) and Leavesly et al., J. Cell Biol., 121:163-170 (1993)).

Briefly, human umbilical vein endothelial cells in log phase and at low passage number were harvested by gentle trypsinization, washed and resuspended at a concentration of 2×10^6 cells/ml in 37°C HBS containing 1% BSA (20 mM HEPES, 150 mM NaCl, 1.8 mM CaCl_2 , 1.8 mM MgCl_2 , 5 mM KCl, and 5 mM glucose, pH 7.4). Antibodies (Vitaxin, LM609, and IgG₁ control) were diluted to 10 µg/ml from stock solutions. Antibodies were added to cells in a 1:1 dilution (final concentration of antibodies = 5 µg/ml; final concentration of cells = 1×10^6 cells/ml) and incubated on ice for 10 - 30 minutes. The cell/antibody suspensions (200 µl to each compartment) were then added to the upper compartments of a Transwell cell culture chamber (Corning Costar), the lower compartments of which had been coated with 0.5 ml of 10 µg/ml vitronectin (in HBS). Vitronectin serves as the chemoattractant for the endothelial cells. The chambers were placed at 37°C for 4 hours to allow cell migration to occur.

Visualization of cell migration was performed by first removing the remaining cells in the upper compartment with a cotton swab. Cells that had migrated to the lower side of the insert were stained with crystal violet for 30 minutes, followed by solubilization in acetic acid and the absorbance of the dye was measured at a wavelength of 550 nm. The amount of absorbance is directly proportional to the number of cells that have migrated from the upper to the lower chamber. The results of the assay are presented in Figure 7B. Both Vitaxin and the parental antibody LM609 yielded essentially identical inhibitory results. Specifically, Vitaxin and LM609 inhibited about 60% of the vitronectin-induced migration of endothelial cells compared to the IgG₁ control and to a sample with no inhibitor.

EXAMPLE IV

Vitaxin-Mediated Inhibition of $\alpha_v\beta_3$ In Animal Models

This Example describes the inhibition of tumor growth by Vitaxin in two animal models. Tumor growth was
 5 inhibited by inhibiting at least $\alpha_v\beta_3$ -mediated neovascularization with Vitaxin.

The first model measures angiogenesis in the chick chorioallantoic membrane (CAM). This assay is a well recognized model for *in vivo* angiogenesis because
 10 the neovascularization of whole tissue is occurring. Specifically, the assay measures growth factor induced angiogenesis of chicken CAM vessels growing toward the growth factor-impregnated filter disk or into the tissue grown on the CAM. Inhibition of neovascularization is
 15 based on the amount and extent of new vessel growth or on the growth inhibition of tissue on the CAM. The assay has been described in detail by others and has been used to measure neovascularization as well as the neovascularization of tumor tissue (Ausprunk et al., Am. J. Pathol., 79:597-618 (1975); Ossonski et al. Cancer Res., 40:2300-2309 (1980); Brooks et al. Science, 264:569-571 (1994a) and Brooks et al. Cell, 79:1157-1164 (1994b)).
 20

Briefly, for growth factor induced angiogenesis
 25 filter disks are punched from #1 Whatman Qualitative Circles using a skin biopsy punch. Disks are first sterilized by exposure to UV light and then saturated with varying concentrations of TNF- α or HBSS as a negative control (for at least 1 hour) under sterile
 30 conditions. Angiogenesis is induced by placing the saturated filter disks on the CAMs.

Inhibition of angiogenesis is performed by treating the embryos with various amounts of Vitaxin and controls (antibody or purified human IgG₁). The treatments are performed by intravenous injection approximately 24 hours after disk placement. After 48 hours, CAMs are dissected and angiogenesis is scored on a scale of 1-4. HBSS saturated filter disks are used as the negative control, representing angiogenesis that may occur in response to tissue injury in preparing CAMs, and, values for these CAMS are subtracted out as background. Purified human IgG₁ is used as the negative control for injections since Vitaxin is of the human IgG₁ subclass. Vitaxin was found to inhibit TNF- α induced angiogenesis in a dose dependent manner. Maximal inhibition occurred with a single dose of Vitaxin at 300 μ g which resulted in greater than 80% inhibition compared to the human IgG₁ control.

In addition to the above described CAM assay using growth factor-induced neovascularization, additional studies were performed utilizing tumor-induced neovascularization. For these assays, angiogenesis was induced by transplanting of $\alpha_v\beta_3$ -negative tumor fragments into the CAMs. The use of $\alpha_v\beta_3$ -negative tumor fragments ensures that any inhibition of tumor growth is due to the inhibition of $\alpha_v\beta_3$ -mediated neovascularization by CAM-derived endothelial cells and not to adhesion events mediated by $\alpha_v\beta_3$ present on the tumor cells.

Inhibition of tumor growth was assessed by placing a single cell suspension of FG (8×10^6 cells, pancreatic carcinoma) and HEP-3 cells (5×10^5 cells, laryngeal carcinoma) onto CAMs in 30 μ l. One week later, tumors are removed and cut into approximately 50 mg fragments at which time they are placed onto new CAMs.

After 24 hours of this second placement embryos are injected intravenously with Vitaxin or human IgG₁ as a negative control. The tumors are allowed to grow for about 7 days following which they are removed and weighed.

The results of Vitaxin treatment on the neovascularization of tumors is shown in Figure 6A. The data is expressed as a mean change in tumor weight and demonstrate that Vitaxin is able to inhibit the growth of $\alpha_v\beta_3$ -negative tumors such as FG and HEP-3 tumor fragments. More specifically, there was a mean weight change for Vitaxin treated FG tumor fragments of -5.38 whereas a change of -11.0 was observed for Vitaxin treated HEP-3 tumors. The IgG₁ controls exhibited positive mean weight changes of 25.29 and 28.5 for the FG and HEP-3 tumor fragments, respectively. These results were obtained following a single intravenous injection.

In a second animal model, the inhibition of Vx2 carcinoma cells in rabbits was used as a measure of Vitaxin's inhibitory effect on tumors. The Vx2 carcinoma is a transplantable carcinoma derived from a Shope virus-induced papilloma. It was first described in 1940 and has since been used extensively in studies on tumor invasion, tumor-host interactions and angiogenesis. The Vx2 carcinoma is fibrotic in nature, highly aggressive, and exhibits features of an anaplastic type carcinoma. Propagation of Vx2 tumor is accomplished through serial transplantation in donor rabbits. Following subcutaneous transplantation, it has been reported that after an initial inflammatory reaction, host repair mechanisms set in between days 2 and 4. This repair mechanism is characterized by the formation of new connective tissue and the production of new capillaries. The newly formed

capillaries are restricted to the repair zone at day 4, however, by day 8 they have extended to the outer region of the tumor. These characteristics and the pharmacokinetics of Vitaxin in rabbits were used to
5 determine initial doses and scheduling of treatments for these experiments. The elimination half life of Vitaxin in animal serum dosed at 1, 5, and 10 mg/kg was found to be 38.9, 60.3, and 52.1 hours, respectively.

Growth of Vx2 tumors in the above animal model
10 was used to study the effect of Vitaxin after early administration on primary tumor growth in rabbits implanted subcutaneously with Vx2 carcinoma. Briefly, Vx2 tumors (50 mg) were transplanted into the inner thigh of rabbits through an incision between the skin and
15 muscle. Measurements of the primary tumor were taken throughout the experiment through day 25. At day 28 after the transplantation animals were sacrificed and tumors were excised and weighed. By day 28, tumors became extremely irregular in shape and as a result,
20 measurements became difficult and were not reflective of tumor volume. Therefore measurements were assessed only through day 25.

In a first study, rabbits were treated starting at day 1 post tumor implantation with 5 and 1 mg/kg
25 Vitaxin every four days for 28 days for a total of 7 doses). In both groups, inhibition of tumor growth was observed. In a second series of studies, rabbits were treated beginning at day 7 post tumor implantation as described above for a total of 5 doses. Inhibition of
30 tumor growth was also observed.

It should be noted that administering a grafted antibody as a repeat dose treatment to rabbits might

generate an immune response that can have a neutralizing effect on Vitaxin thus potentially comprising efficacy. Preliminary data suggest that approximately 25-50% of the animals develop such a response.

5 The results of each of the Vitaxin treatments described above is shown in Figure 6B and 6C. In the rabbits receiving treatments on day 1, inhibition of tumor growth was observed in both the 1 mg/kg and the 5 mg/kg dosing groups compared to the control PBS treated
10 control. Specifically, a growth inhibition of about 67 and 80% was observed, respectively, as measured by the mean tumor weight. A lesser degree of inhibition was observed in animals that began Vitaxin treatment on day 7 post implantation. These results are shown in Figure 6C.
15 In all cases, inhibition of tumor growth was not seen at Vitaxin concentrations lower than 0.2 mg/kg.

EXAMPLE V

Construction of LM609 Grafted Functional Antibody Fragments

20 This Example shows the construction of functional LM609 grafted antibody fragments in which only the CDRs have been transferred from the LM609 donor antibody to a human acceptor framework.

25 CDR grafting of LM609 to produce a functional antibody fragment was accomplished by the methods set forth below. These procedures are applicable for the CDR grafting of essentially any donor antibody where amino acid residues outside of the CDRs from the donor antibody are not desired in the final grafted product.

5.0). Human framework sequences showing significant identity to the murine LM609 variable region gene sequences were selected for receiving the LM609 CDRs. Human heavy chain variable region M72 'CL had 88% identity to frameworks 1, 2 and 3 of LM609 heavy chain and human light chain V region LS1 'CL had 79% identity to frameworks 1, 2 and 3 of LM609 light chain. With the exclusion of non-identical residues outside of the CDRs as defined by Kabat et al. murine LM609 CDR sequences as defined by Kabat et al. and MacCallum et al. were grafted onto the human frameworks. Using this grafting scheme, the final grafted product does not contain any amino acid residues outside of the CDRs as defined by Kabat et al. which are identical to an LM609 amino acid at the corresponding position (outside of residues: 31-35, 50-66 and 99-106 for CDRs 1, 2 and 3, respectively, of the heavy chain variable region and residues 24-34, 50-56, and 89-97 for CDRs 1, 2 and 3, respectively, of the light chain variable region). Moreover, no intermediates are produced which contain an amino acid residue outside of the CDRs as defined by Kabat et al. which are identical to the LM609 amino acid at that position. The CDR grafting procedures are set forth below.

Full-length CDR grafted variable region genes were synthesized by PCR using long overlapping oligonucleotides as described previously in Example II. The heavy chain variable region oligonucleotides were those described previously as SEQ ID NOS:9-13. The light chain variable region oligonucleotides were synthesized so as to contain the CDR grafted variable region as well as a stop codon at position 49. The five oligonucleotides for the light chain LM609 grafted variable region are show as SEQ ID NOS:23-27 where the second oligonucleotide in the series contains the stop

codon at position 49 (SEQ ID NO:24). The nucleotide sequences of oligonucleotides used to construct LM609 grafted light chain variable region is shown in Table 7.

Table 7: Oligonucleotides Used to Construct LM609

5 **Grafted Light Chain Variable Region**

GAGATTGTGC TAACTCAGTC TCCAGCCACC CTGTCTCTCA GCCCAGGAGA AAGGGCGACT
 CTTTCCTGCC AGGCCAGCCA AAGTATT SEQ ID NO: 23
 TTAGATGAGA AGCCTTGGGG CTTGACCAGG CCTTTGTTGA TACCAGTGTA GGTGGTTGCT
 AATACTTTGG CTGGC SEQ ID NO: 24
 10 CCAAGGCTTC TCATCTAATA TCGTTCCCAG TCCATCTCTG GGATCCCCGC CAGGTTTCAGT
 GGCAGTGGAT CAGGGACAGA TTTC SEQ ID NO: 25
 GCTGCCACTC TGTTGACAGT AATAGACTGC AAAATCTTCA GGCTCCAGAC TGGAGATAGT
 GAGGGTGAAA TCTGTCCCTG A SEQ ID NO: 26
 CAACAGAGTG GCAGCTGGCC TCACACGTTC GGAGGGGGGA CCAAGGTGGA AATTAAG
 15 SEQ ID NO:27

All long oligonucleotides were gel purified. CDR grafting of the LM609 heavy chain variable region was constructed by mixing 5 overlapping oligonucleotides (SEQ ID NOS:9-13), at equimolar concentrations, in the
 20 presence of annealing PCR primers containing at least 18 nucleotide residues complementary to vector sequences for the efficient annealing of the amplified V region product to the single-stranded vector. The annealed mixture was fully converted to a double-stranded molecule with T4 DNA
 25 polymerase plus dNTPs and ligated with T4 ligase. The mutagenesis reaction (1 μ l) was electroporated into *E. coli* strain DH10B (BRL), titered onto a lawn of XL-1 (Stratagene, Inc.) and incubated until plaques formed. Replica filter lifts were prepared and plaques containing
 30 V_H gene sequences were screened either by hybridization with a digoxigenin-labeled oligonucleotide complementary to LM609 heavy chain CDR 2 sequences or reactivity with 7F11-alkaline phosphatase conjugate, a monoclonal

BB06707909T060

antibody raised against the decapeptide sequence Tyr Pro Tyr Asp Val Pro Asp Tyr Ala Ser (SEQ ID NO:28) appended to the carboxy terminus of the vector CH₁ domain (Biosite, Inc., San Diego, CA).

5 Fifty clones that were double-positive were pooled and used to prepare uridinylated template for hybridization mutagenesis with the amplified CDR grafted LM609 V_L product constructed in a similar fashion using the five overlapping oligonucleotides shown as SEQ ID
10 NOS:23-27. The mutagenesis reaction was electroporated into *E. coli* strain DH10B. Randomly picked clones were sequenced to identify a properly constructed template for construction of the non-donor library at position 49. This template was prepared as a uridinylated template and
15 an oligonucleotide population of the following sequence was used for site directed mutagenesis.

GGGAACGATA-19aa-GATGAGAAGC

The sequence 19aa in the above primer (SEQ ID NO:30) represents the fact that this primer specifies a sequence
20 population consisting of 19 different codon sequences that encode each of the 19 non-donor amino acids. These amino acids are those not found at position 49 of LM609 and include all amino acids except for Lys. Clones that resulted from this mutagenesis were picked and antibody
25 expressed by these clones were prepared. These samples were then screened for binding to $\alpha\text{v}\beta_3$ in an ELISA assay. Clones having either Arg or Met amino acids in position 49 were functionally identified. The nucleotide and amino acid sequence of the LM609 grafted heavy chain
30 variable region is show in Figure 1A (SEQ ID NOS:1 and 2, respectively). The nucleotide and amino acid sequence of

860570 "T909T060

the LM609 grafted light chain variable region is shown in Figure 7 (SEQ ID NOS:31 and 32, respectively).

EXAMPLE VI

Generation of LM609 Grafted Antibodies Having Enhanced Activity

5

This example shows *in vitro* maturation of LM609 grafted antibody to obtain antibody variants having increased affinity to $\alpha_v\beta_3$ relative to the parent LM609 grafted antibody.

10 To optimize the affinity of LM609 grafted antibody *in vitro*, an M13 phage system was used, which permits the efficient synthesis, expression, and screening of libraries of functional antibody fragments (Fabs). The contribution of each of the six CDRs of the
15 Ig heavy and light chains was assessed. The CDRs were defined broadly based on a combination of sequence variability and antibody structural models (Kabat et al., J. Biol. Chem. 252:6609-6616 (1977); Chothia et al., *supra*; MacCallum et al., *supra*). Thus, one library was
20 constructed for each CDR, with the exception of H2 which was split into two libraries due to its long (20 amino acids) length. The variable region frameworks which harbored the mutated CDRs were the heavy chain variable region shown in Figure 1a (SEQ ID NO:2) and the light
25 chain variable region shown in Figure 7 (SEQ ID NO:32).

CDRs were chosen from the heavy chain variable region shown in Figure 1a (SEQ ID NO:2) and the light chain variable region shown in Figure 7 (SEQ ID NO:32). Briefly, utilizing the numbering system of Kabat et al.,
30 *supra*, the residues chosen for mutagenesis of the CDRs

"T909T060"

(Table 9) were: Gln²⁴-Tyr³⁶ in light chain CDR1 (L1);
 Leu⁴⁶-Ser⁵⁶ in light chain CDR2 (L2); Gln⁸⁹-Thr⁹⁷ in light
 chain CDR3 (L3); Gly²⁶-Ser³⁵ in heavy chain CDR1 (H1);
 Trp⁴⁷-Gly⁶⁵ in heavy chain CDR2 (H2); and Ala⁹³-Tyr¹⁰² in
 5 heavy chain CDR3 (H3). Libraries were created for each
 CDR, with the oligonucleotides designed to mutate a
 single CDR residue in each clone. Due to the extended
 length of H2, two libraries mutating residues 47-55 (H2a)
 and 56-65 (H2b), respectively, were constructed to cover
 10 this region.

The template for generating light chain CDR3
 mutants contained Gly at position 92. However, it was
 subsequently determined that position 92 of the light
 chain CDR3 was inadvertently deduced to be a Gly,
 15 resulting in humanized LM609 grafted antibodies being
 constructed with Gly at that position. It was later
 realized that the original LM609 sequence contained an
 Asn at position 92. Using the methods described herein
 to introduce mutations into CDRs of an LM609 grafted
 20 antibody, an LM609 grafted antibody having Asn at
 position 92 of light chain CDR3 was found to have $\alpha_v\beta_3$
 binding activity (see Table 9), confirming the
 identification of Asn⁹² as a functional LM609 grafted
 antibody. Thus, antibodies containing light chain CDR3
 25 having Gly or Asn at position 92 are active in binding
 $\alpha_v\beta_3$.

Oligonucleotides encoding a single mutation
 were synthesized by introducing NN(G/T) at each CDR
 position as described previously (Glaser et al., *supra*).
 30 The antibody libraries were constructed in M131XL604
 vector by hybridization mutagenesis as described
 previously, with some modifications (Rosok et al., J.
Biol. Chem. 271:22611-22618 (1996); Huse et al., J.

Variants that bound $\alpha_v\beta_3$ were identified by incubating the filters for 10-15 min in 0.1 M Tris, pH 9.5, containing 0.4 mM 2,2'-di-*p*-nitrophenyl-5,5'-diphenyl-3,3'-(3,3'-dimethoxy-4,4'-diphenylene)ditetrazolium chloride and 0.38 mM 5-bromo-4-chloro-3-indoxyl phosphate mono-(*p*-toluidinium) salt (JBL Scientific, Inc.; San Luis Obispo, CA).

To generate biotinylated $\alpha_v\beta_3$, the $\alpha_v\beta_3$ receptor was purified from human placenta by affinity chromatography, as described previously (Smith and Cheresch, J. Biol. Chem. 263:18726-18731 (1988)). To biotinylate $\alpha_v\beta_3$, purified receptor was dialyzed into 50 mM HEPES, pH 7.4, 150 mM NaCl, 1.0 mM CaCl₂, containing 0.1% NP-40 (binding buffer) and incubated with 100-fold molar excess sulfosuccinimidobiotin for 3h at 4°C. The reaction was terminated by the addition of 50 mM ethanolamine.

Phage expressed LM609 grafted antibody variants were selectively captured on nitrocellulose filters coated with goat anti-human kappa chain antibody, probed with biotinylated $\alpha_v\beta_3$, and detected with NeutrAvidin-alkaline phosphatase. Initially, biotinylated $\alpha_v\beta_3$ was titrated on lifts containing phage expressing the LM609 grafted antibody parent molecule only. Subsequently, the concentration of biotinylated $\alpha_v\beta_3$ was decreased to yield a barely perceptible signal. In this way, only clones expressing higher affinity variants were readily identified during screening of the variant libraries. Following the exhaustive capture lift screening of ≥ 2500 clones from each library, 300 higher affinity variants were identified (see Table 10). The greatest number of clones displaying improved affinity were identified in the H3 (185) and L3 (52) CDRs, though

variants with improved affinity were identified in every CDR.

LM609 grafted antibody variants identified by capture lift as having $\alpha_v\beta_3$ binding activity were further characterized to determine binding affinity to $\alpha_v\beta_3$, specificity for $\alpha_v\beta_3$ over other integrins, and $\alpha_v\beta_3$ association and dissociation rates. For these assays, purified Fab of LM609 grafted antibody variants was used. Briefly, Fab was expressed as described previously and was released from the periplasmic space by sonic oscillation (Watkins et al., *supra*, 1997). Cells collected from one liter cultures were lysed in 10 ml 50 mM Tris, pH 8.0, containing 0.05% Tween 20. Fab was bound to a 1 ml protein A column (Pharmacia) which had been equilibrated with 50 mM glycine, pH 8, containing 250 mM NaCl, washed with the same buffer, and eluted with 10 ml of 100 mM glycine, pH 3, into one-tenth volume 1 M Tris, pH 8. Purified Fab was quantitated as described previously (Watkins et al., *supra*, 1997).

LM609 grafted antibody variants were tested for binding to $\alpha_v\beta_3$ and specificity of binding to $\alpha_v\beta_3$ relative to $\alpha_v\beta_5$ and $\alpha_{Iib}\beta_3$. For ELISA titration of Fab on immobilized $\alpha_v\beta_3$ and the related integrins $\alpha_v\beta_5$ and $\alpha_{Iib}\beta_3$, Immulon II microtiter plates were coated with 1 μ g/ml purified receptor in 20 mM Tris, pH 7.4, 150 mM NaCl, 2 mM CaCl_2 , 1 mM MgCl_2 , 1 mM MnCl_2 , washed once, and blocked in 3% BSA in 50 mM Tris, pH 7.4, 100 mM NaCl, 2 mM CaCl_2 , 1 mM MgCl_2 for 1 h at 25°C. Human $\alpha_{Iib}\beta_3$, purified from platelets, was obtained from Enzyme Research Laboratories, Inc. (South Bend, IN) and $\alpha_v\beta_5$ was purified from placental extract depleted of $\alpha_v\beta_3$, as described previously (Smith et al., *J. Biol. Chem.* 265:11008-11013 (1990)). Just prior to use, the plates were washed two

times and were then incubated 1 h at 25°C with various dilutions of Fab. The plates were washed five times, incubated 1 h at 25°C with goat anti-human kappa-alkaline phosphatase diluted 2000-fold, washed five times, and developed as described previously (Watkins et al., *supra*, 1997). All dilutions and washes were in 50 mM Tris-HCl, pH 7.4, 100 mM NaCl, 2 mM CaCl₂, and 1 mM MgCl₂.

Table 10: Capture Lift Screening of LM609 grafted antibody CDR Libraries.

Library	Size ¹	Screened ²	Positives ³	Enhanced Affinity ⁴
H1	320	2500	16	8
H2a	320	5000	26	7
H2b	320	5000	2	1
H3	320	5000	185	78 ⁵
L1	416	2500	12	1
L2	352	3250	7	1
L3	288	5000	52	41

¹Number of unique clones based on DNA sequence. Thirty-two condons are used to express all twenty amino acids at each position.

²Phage-expressed libraries were plated on XL-1 Blue/agar lawns at 500-100 plaques per 100 mm dish.

³Positives are defined as clones that were identified in the initial screen, replated, and verified in a second capture lift assay.

⁴Soluble Fab was titrated against immobilized $\alpha_v\beta_3$ in an ELISA format. Based on comparison of the inflection

point of the titration profiles, clones which displayed ~3-fold enhanced affinity were selected for further characterization.

⁵Of the 185 positive clones identified by capture lift, 98 were further characterized for binding to immobilized $\alpha_V\beta_3$.

Figure 8 shows titration of antibody variants and LM609 grafted antibody Fab on immobilized $\alpha_V\beta_3$. Bacterial cell lysates containing LM609 grafted antibody (closed circles), variants with improved affinity isolated from the primary libraries (S102, closed squares; Y100, open squares; and Y101, open triangles) or from the combinatorial libraries (closed triangles), or an irrelevant Fab (open circles) were titrated on immobilized $\alpha_V\beta_3$.

Comparison of the inflection points of the binding profiles obtained from titrating variants on immobilized $\alpha_V\beta_3$ demonstrated that multiple clones displayed >3-fold improved affinity, confirming the effectiveness of utilizing the capture lift in a semi-quantitative fashion (Figure 8, compare squares and open triangles with closed circles). Based on the capture lift screening and subsequent characterization of binding to immobilized $\alpha_V\beta_3$, it was concluded that both heavy and light chain CDRs are directly involved in the interaction of $\alpha_V\beta_3$ with the LM609 grafted antibody variants.

DNA was isolated from clones displaying >3-fold enhanced binding and sequenced to identify the mutations which resulted in higher affinity. DNA sequencing was performed on isolated single-stranded DNA. The heavy and light chain variable region genes were sequenced by the

fluorescent dideoxynucleotide termination method (Perkin-Elmer; Foster City, CA). Based on sequence analysis of 103 variants, 23 unique mutations clustered at 14 sites were identified (Table 9). The majority of the sites of beneficial mutations were found in the heavy chain CDRs, with four located in H3, and three each in H2 (2a and 2b combined) and H1. Seven distinct and beneficial amino acid substitutions were identified at a single site within H3, tyrosine residue 102. The diverse nature of the substitutions at this site suggests that tyrosine residue 102 may sterically hinder LM609 grafted antibody binding to $\alpha_v\beta_3$. In support of this, variants expressing the other aromatic amino acids (phenylalanine, histidine, and tryptophan) instead of tyrosine at residue 102 were never isolated following screening for enhanced binding.

The affinities of select variants were further characterized by utilizing surface plasmon resonance (BIAcore) to measure the association and dissociation rates of purified Fab with immobilized $\alpha_v\beta_3$. Briefly, surface plasmon resonance (BIAcore; Pharmacia) was used to determine the kinetic constants for the interaction between $\alpha_v\beta_3$ and LM609 grafted antibody variants. Purified $\alpha_v\beta_3$ receptor was immobilized to a (1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide hydrochloride)/N-hydroxysuccinimide-activated sensor chip by injecting 30 μ l of 15 μ g/ml $\alpha_v\beta_3$ in 10 mM sodium acetate, pH 4. To obtain association rate constants (k_{on}), the binding rate at five different Fab concentrations, ranging from 5-40 μ g/ml in 50 mM Tris-HCl, pH 7.4, 100 mM NaCl, 2 mM $CaCl_2$, and 1 mM $MgCl_2$, was determined at a flow rate of 10 μ l/min. Dissociation rate constants (k_{off}) were the average of five measurements obtained by analyzing the dissociation phase

at an increased flow rate (40 $\mu\text{l}/\text{min}$). Sensorgrams were analyzed with the BIAevaluation 2.1 program (Pharmacia). Residual Fab was removed after each measurement with 10 mM HCl, 2 mM CaCl_2 and 1 mM MgCl_2 .

5 Table 9 shows that the variants all displayed a lower K_d than the LM609 grafted antibody parent molecule, consistent with both the capture lift and the ELISA. Analysis of association and dissociation rates revealed that the majority of improved variants had slower
10 dissociation rates while having similar association rates. For example, LM609 grafted antibody had an association rate $18.0 \times 10^4 \text{ M}^{-1}\text{s}^{-1}$, while the variants ranged from $16.7\text{--}31.8 \times 10^4 \text{ M}^{-1}\text{s}^{-1}$. In contrast, every clone dissociated slower than LM609 grafted antibody
15 ($4.97 \times 10^{-3} \text{ s}^{-1}$) with dissociation rates ranging from 1.6-fold ($3.03 \times 10^{-3} \text{ s}^{-1}$) to 11.8-fold ($0.42 \times 10^{-3} \text{ s}^{-1}$) slower.

 These results demonstrate that introducing single amino acid substitutions into LM609 grafted
20 antibody CDRs allows the identification of modified LM609 grafted antibodies having higher affinity for $\alpha_v\beta_3$ than the parent LM609 grafted antibody.

Table 9: Identification of Enhanced IM609 Grafted Antibodies from Primary Libraries

chain*	library†	sequence	k_{on} ($\times 10^4$) ($M^{-1}s^{-1}$)	k_{off} ($\times 10^{-3}$) (s^{-1})	Kd (nM)
	LM609 grafted antibody		18.0	4.97	27.6
H	CDR1 T27 W29 L30	G F T F S S Y D M S T W L	n.d. n.d. n.d.	n.d. n.d. n.d.	n.d. n.d. n.d.
H	CDR2a K52	W V A K V S S G G G K	17.8	2.18	12.2
H	CDR2b P60 E64	S T Y Y L D T V Q G P E	31.8 n.d.	1.85 n.d.	5.8 n.d.
H	CDR3 H97 Y100 D101 Y101 S102 T102 D102 E102 M102 G102 A102	A R H N Y G S F A Y H Y Y D Y S T D E M G A	22.0 17.5 n.d. 21.8 24.2 24.6 27.6 n.d. n.d. 16.1 27.5	3.03 2.51 n.d. 0.48 1.44 1.43 0.97 n.d. n.d. 2.01 2.27	13.8 14.3 n.d. 2.2 6.0 5.8 3.5 n.d. n.d. 12.5 8.3
L	CDR1 F32	Q A S Q S I S N H L H W Y F	16.7	0.42	2.5
L	CDR2 S51	L L I R Y R S Q S I S S	n.d.	n.d.	n.d.
L	CDR3 N92 T92 L96 Q96	Q Q S G S W P H T N T L Q	23.6 n.d. 24.3 n.d.	1.35 n.d. 2.23 n.d.	5.7 n.d. 9.2 n.d.

EXAMPLE VIIGeneration of High Affinity LM609 Grafted Antibodies

This example shows that single amino acid mutations in CDRs of an LM609 grafted that result in
5 higher affinity binding to $\alpha_v\beta_3$ can be combined to generate high affinity LM609 grafted antibodies.

Random combination of all of the beneficial mutations of LM609 grafted antibody would generate a combinatorial library containing $>10^5$ variants, requiring
10 efficient screening methodologies. Therefore, to determine if clones displaying >10 -fold enhanced affinities could be rapidly distinguished from one another, variants displaying 3 to 13-fold enhanced affinity were evaluated by capture lift utilizing lower
15 concentrations of biotinylated $\alpha_v\beta_3$. Despite repeated attempts with a broad range of concentrations of $\alpha_v\beta_3$, consistent differences in the capture lift signals were not observed. Because of this, smaller combinatorial libraries were constructed and subsequently screened by
20 ELISA.

Four distinct combinatorial libraries were constructed in order to evaluate the optimal number of combinations that could be accomplished utilizing two site hybridization mutagenesis (Figure 9). Briefly,
25 combinatorial libraries were constructed by synthesizing degenerate oligonucleotides encoding both the wild-type and beneficial heavy chain mutations (H2, Leu⁶⁰→Pro; H3 Tyr⁹⁷→His; H3, Ala¹⁰¹→Tyr; H3, Tyr¹⁰²→Ser, Thr, Asp, Glu, Met, Gly, Ala). Utilizing two site hybridization
30 mutagenesis, as described above, the oligonucleotides were annealed at a 40:1 molar ratio to uridinylated template prepared from LM609 grafted antibody and three

light chain mutations (Figure 9; L1, His³²→Phe; L3, Gly⁹²→Asn; L3, His⁹⁶→Leu). As a result, a total of 256 variants were synthesized in four combinatorial library subsets.

5 Figure 9 shows construction of combinatorial libraries of beneficial mutations. Uridinylated template from LM609 grafted antibody and three optimal light chain variants (F32, N92, and L96) was prepared. Two site hybridization was performed with two degenerate
10 oligonucleotides, which were designed to introduce beneficial mutations at four distinct heavy chain residues.

 Following preparation of uridinylated templates of LM609 grafted antibody and three light chain variants,
15 (Table 9; F32, N92, and L96), degenerate oligonucleotides encoding the wild type residue and the most beneficial heavy chain mutations (Table 9; P60, H97, Y101, S102, T102, D102, E102, M102, G102, and A102) were hybridized to the light chain templates, resulting in four
20 combinatorial libraries, each containing 64 unique variants. Potentially, the combination of multiple mutations can have detrimental effects on affinity and, thus, can prevent the identification of beneficial combinations resulting from mutations at fewer sites.
25 For this reason, the amino acid expressed by the LM609 grafted antibody parent molecule was included at each position in the combinatorial library. By utilizing this approach, simultaneous combinatorial mutagenesis of three CDRs (L1 or L3 each in combination with H2 and H3) was
30 accomplished. Based on sequence analysis, the two site hybridization mutagenesis was achieved with ~50% efficiency.

In order to screen the combinatorial libraries, soluble Fab was expressed and released from the periplasm of small-scale (<1 ml) bacterial cultures that had been infected with randomly selected clones. Although

5 variable expression levels were observed, uniform quantities of the unpurified variants were captured on a microtiter plate through a peptide tag present on the carboxyl-terminus of the heavy chain. Briefly, combinatorial LM609 grafted antibody libraries were

10 screened by an ELISA that permits the determination of relative affinities of antibody variants produced in small-scale bacterial cultures (Watkins et al., Anal. Biochem. 253:37-45 (1997)). An Immulon II microtiter plate (Dynatech Laboratories; Chantilly, VA) was coated

15 with 10 $\mu\text{g/ml}$ of the 7F11 monoclonal antibody, which recognizes a peptide tag on the carboxyl-terminus of the LM609 grafted antibody variant heavy chains (Field et al., Mol. Cell. Biol. 8:2159-2165 (1988)). Following capture of Fab from *E. coli* lysates, the plate was

20 incubated with 0.5-1 $\mu\text{g/ml}$ biotinylated $\alpha_v\beta_3$ for 1 h at 25°C. The plate was washed seven times, incubated with 0.5 U/ml streptavidin-alkaline phosphatase (1000 U/ml; Boehringer Mannheim; Indianapolis, IN) for 15 min at 25°C, washed seven times, and developed as described

25 previously (Watkins et al., *supra*, 1997). All dilutions and washes were in binding buffer.

As described previously (Watkins et al., *supra*, 1997), this ELISA screening method enabled a rapid and direct comparison of the relative affinities of the

30 variants following incubation with biotinylated $\alpha_v\beta_3$ and streptavidin-alkaline phosphatase. To ensure that the full Fab diversity was sampled, one thousand randomly selected clones were screened from each combinatorial library. Variants that displayed an enhanced ELISA

signal were further characterized for binding to immobilized $\alpha_v\beta_3$ (Figure 8, closed triangles) and were sequenced to identify the mutations (Table 10).

Screening of the four combinatorial libraries identified fourteen unique combinations of mutations that improved binding significantly over the individual mutations identified in the screening of the first library. While the best clone from the primary screen had a 12.5-fold increase in affinity, the fourteen unique combinations isolated from screening the combinatorial libraries displayed affinities ranging from 18 to 92-fold greater than the parent LM609 grafted antibody. The majority of these variants consisted of H2 and H3 mutations combined with the L1 or L3 mutations. Beneficial combinations of heavy chain mutations with wild-type light chain were also identified, but did not result in improved affinity to the same extent as other combinatorial variants. The variants predominantly contained 2 to 4 mutations, with one clone, C29, containing five mutations. No direct correlation between the total number of mutations in each variant and the resulting affinity was observed. For example, while the binding of clone C37 was 92-fold enhanced over the parent molecule and was achieved through the combination of three mutations, clone C29 had ~55-fold greater affinity achieved through the combination of five mutations. Multiple variants displaying >50-fold enhanced affinity resulting from the combination of as few as two mutations were identified (2G4, 17, and V357D).

The combinatorial clones with improved affinity all displayed >10-fold slower dissociation rates, possibly reflecting a selection bias introduced by long incubation steps in the screening. In addition, all of

09016061-013098

5
10

15

Table 10: Identification of Optimal Combinatorial Mutations

Library*	clone	L1 32	L3 92	L3 96	H2 60	H3 97	H3 101	H3 102	k_{on} ($\times 10^4$) ($M^{-1}s^{-1}$)	k_{on} ($\times 10^{-3}$) (s^{-1})	Kd (nM)
wild type		H	G	H	L	Y	A	Y	18.0	4.97	27.6
F32	17	F						S	25.1	0.138	0.5
	7	F			P	H		S	20.4	0.236	1.2
	56	F			P			S	26.6	0.135	0.5
	C59	F			P			D	26.5	0.137	0.5
	C176	F			P			T	22.5	0.192	0.9
	V357D	F						D	27.9	0.140	0.5
N92	C119		N		P			S	21.5	0.316	1.5
L96	8F9			L	P	H		S	47.5	0.280	0.6
	C29			L	P	H	Y	S	67.5	0.343	0.5
	2G4			L				S	60.3	0.229	0.4
	6H6			L		H		S	50.4	0.187	0.4
	C37			L			Y	E	44.8	0.147	0.3
	6D1			L	P		Y	S	41.0	0.158	0.4
	6G1			L	P			S	38.9	0.280	0.7

As a first step toward determining if the increase in affinity of the variants resulted in greater biological activity, variants displaying a range of affinities were assayed for their ability to inhibit the binding of a natural ligand, fibrinogen, to immobilized $\alpha_v\beta_3$ receptor. Briefly, LM609 grafted antibody variants were tested for inhibition of ligand binding as described previously except that the binding of biotinylated human fibrinogen (Calbiochem, La Jolla, CA) was detected with 0.5 $\mu\text{g/ml}$ NeutrAvidin-alkaline phosphatase (Smith et al., J. Biol. Chem. 265:12267-12271 (1990)).

The results of these competition assays are shown in Figure 10. Figure 10A shows inhibition of fibrinogen binding to immobilized $\alpha_v\beta_3$. Immobilized $\alpha_v\beta_3$ was incubated with 0.1 $\mu\text{g/ml}$ biotinylated fibrinogen and various concentrations of LM609 grafted antibody (open circles), S102 (closed circles), F32 (open triangles), or C59 (closed triangles) for 3 h at 37°C. Unbound ligand and Fab were removed by washing and bound fibrinogen was quantitated following incubation with NeutrAvidin alkaline phosphatase conjugate. Figure 10B shows correlation of affinity of variants with inhibition of fibrinogen binding. The concentration of variants required to inhibit the binding of fibrinogen to immobilized $\alpha_v\beta_3$ by 50% (IC_{50}) was plotted as a function of the affinity (Kd).

As shown in Figure 10A, higher affinity variants were more effective at blocking the ligand binding site of the receptor (compare LM609 grafted antibody, open circles, with any of the variants). Subsequent analysis of ten variants displaying affinities (Kd) ranging from 0.3 to 27 nm demonstrated a good correlation ($r^2 = 0.976$) between affinity and ability to

inhibit fibrinogen binding (Figure 10B). In addition, the variants were tested for inhibition of vitronectin binding to the receptor. Similar to fibrinogen, the variants were more effective at inhibiting the
 5 interaction than the parent molecule. Thus, consistent with the cross-reactivity studies with related integrin receptors, mutations which increased affinity did not appear to substantially alter the manner in which the antibody interacted with the receptor.

10 The ability of the variants to inhibit the adhesion of M21 human melanoma cells expressing the $\alpha_v\beta_3$ receptor to fibrinogen was examined. Inhibition of the adhesion of 4×10^4 M21 cells to fibrinogen by the LM609 grafted antibody variants was performed as described
 15 previously (Leavesley et al., J. Cell Biol. 117:1101-1107 (1992)). Similar to the ligand competition studies with purified fibrinogen and $\alpha_v\beta_3$ receptor, higher affinity variants were generally more effective at preventing cell adhesion than was LM609 grafted antibody (Figure 11).
 20 Figure 11 shows inhibition of M21 human melanoma cell adhesion to fibrinogen. Cells and various concentrations of LM609 grafted antibody Fab (closed triangles), S102 (open circles), G102 (closed circles), or C37 (open triangles) were added to 96 well cell culture plates
 25 which had been coated with 10 $\mu\text{g/ml}$ fibrinogen. After incubating for 35 min at 37°C, unbound cells were removed by washing and adherent cells were quantitated by crystal violet staining.

Although intact LM609 grafted antibody Ig
 30 inhibits cell adhesion, the phage expressed Fab did not affect cell adhesion at concentrations as high as 1 mg/ml (Figure 11, closed triangles). Clone C37, isolated from the combinatorial library and displaying ~90-fold greater

affinity than LM609 grafted antibody Fab, inhibited cell adhesion completely (Figure 11, open triangles). Variant G102 had a moderately higher affinity (2.2-fold enhanced) and also inhibited cell adhesion, though less effectively than C37 (Figure 11, closed circles). Surprisingly, clone S102 (Figure 11, open circles), which had a 4.6-fold higher affinity than LM609 grafted antibody, was ineffective at inhibiting cell adhesion, suggesting that clones G102 and S102 interact with the $\alpha_v\beta_3$ receptor differently.

These results show that combining single amino acid mutations that result in LM609 grafted antibodies exhibiting higher binding affinity to $\alpha_v\beta_3$ allows the identification of high affinity LM609 grafted antibody mutants having greater than 90-fold higher binding affinity than the parent LM609 grafted antibody.

Although the invention has been described with reference to the disclosed embodiments, those skilled in the art will readily appreciate that the specific experiments detailed are only illustrative of the invention. It should be understood that various modifications can be made without departing from the spirit of the invention. Accordingly, the invention is limited only by the following claims.

What is claimed is:

1. A Vitaxin antibody exhibiting selective binding affinity to $\alpha_v\beta_3$, comprising at least one Vitaxin heavy chain polypeptide comprising substantially the same variable region amino acid sequence as that shown in Figure 1A (SEQ ID NO:2) and at least one Vitaxin light chain polypeptide comprising substantially the same variable region amino acid sequence as that shown in Figure 1B (SEQ ID NO:4) or a functional fragment thereof.
2. The Vitaxin antibody of claim 1, wherein said functional fragment is selected from the group consisting of Fv, Fab, F(ab)₂ and scFV.
3. A nucleic acid encoding a Vitaxin heavy chain polypeptide comprising substantially the same Vitaxin heavy chain variable region nucleotide sequences as that shown in Figure 1A (SEQ ID NO:1) or a fragment thereof.
4. The nucleic acid of claim 3, wherein said fragment further comprises a nucleic acid encoding substantially the same nucleotide sequence as the variable region of said Vitaxin heavy chain polypeptide (SEQ ID NO:1).
5. The nucleic acid of claim 3, wherein said fragment further comprises a nucleic acid encoding substantially the same nucleotide sequence as a CDR of said Vitaxin heavy chain polypeptide.

6. A nucleic acid encoding a Vitaxin light chain polypeptide comprising substantially the same Vitaxin light chain variable region nucleotide sequences as that shown in Figure 1B (SEQ ID NO:3) or a fragment thereof.

7. The nucleic acid of claim 6, wherein said fragment further comprises a nucleic acid encoding substantially the same nucleotide sequence as the variable region of said Vitaxin light chain polypeptide (SEQ ID NO:3).

8. The nucleic acid of claim 6, wherein said fragment further comprises a nucleic acid encoding substantially the same nucleotide sequence as a CDR of said Vitaxin light chain polypeptide.

9. A nucleic acid encoding a Vitaxin heavy chain polypeptide comprising a nucleotide sequence encoding substantially the same Vitaxin heavy chain variable region amino acid sequence as that shown in Figure 1A (SEQ ID NO:2) or fragment thereof.

10. The nucleic acid of claim 9, wherein said fragment further comprises a nucleic acid encoding substantially the same heavy chain variable region amino acid sequence of said Vitaxin heavy chain amino acid sequence (SEQ ID NO:2).

11. The nucleic acid of claim 9, wherein said fragment further comprises a nucleic acid encoding substantially the same heavy chain CDR amino acid sequence of said Vitaxin heavy chain amino acid sequence.

12. A nucleic acid encoding a Vitaxin light chain polypeptide comprising a nucleotide sequence encoding substantially the same Vitaxin light chain variable region amino acid sequence as that shown in Figure 1B (SEQ ID NO:4) or fragment thereof.

13. The nucleic acid of claim 12, wherein said fragment further comprises a nucleic acid encoding substantially the same light chain variable region amino acid sequence of said Vitaxin light chain amino acid sequence (SEQ ID NO:4).

14. The nucleic acid of claim 12, wherein said fragment further comprises a nucleic acid encoding substantially the same light chain CDR amino acid sequence of said Vitaxin light chain amino acid sequence.

15. A Vitaxin heavy chain polypeptide comprising substantially the same variable region amino acid sequence as that shown in Figure 1A (SEQ ID NO:2) or functional fragment thereof.

16. The Vitaxin heavy chain polypeptide of claim 15, wherein said functional fragment comprises a variable chain polypeptide or a CDR polypeptide.

17. A Vitaxin light chain polypeptide comprising substantially the same variable region amino acid sequence as that shown in Figure 1B (SEQ ID NO:4) or a functional fragment thereof.

18. The Vitaxin light chain polypeptide of claim 17, wherein said functional fragment comprises a variable chain polypeptide or a CDR polypeptide.

0301661.013098

19. A LM609 grafted antibody exhibiting selective binding affinity to $\alpha_v\beta_3$, comprising at least one LM609 grafted heavy chain polypeptide comprising substantially the same variable region amino acid
5 sequence as that shown in Figure 1A (SEQ ID NO:2) and at least one LM609 grafted light chain polypeptide comprising substantially the same variable region amino acid sequence as that shown in Figure 7 (SEQ ID NO:32) or a functional fragment thereof.

10 20. The LM609 grafted antibody of claim 19, wherein said functional fragment is selected from the group consisting of Fv, Fab, F(ab)₂ and scFV.

21. A nucleic acid encoding a LM609 grafted heavy chain polypeptide comprising substantially the same
15 LM609 grafted heavy chain variable region nucleotide sequences as that shown in Figure 1A (SEQ ID NO:1) or a fragment thereof.

22. The nucleic acid of claim 21, wherein said fragment further comprises a nucleic acid encoding
20 substantially the same nucleotide sequence as the variable region of said LM609 grafted heavy chain polypeptide (SEQ ID NO:1).

23. The nucleic acid of claim 21, wherein said fragment further comprises a nucleic acid encoding
25 substantially the same nucleotide sequence as a CDR of said LM609 grafted heavy chain polypeptide.

SECRET - 013099

24. A nucleic acid encoding a LM609 grafted light chain polypeptide comprising substantially the same LM609 grafted light chain variable region nucleotide sequences as that shown in Figure 7 (SEQ ID NO:31) or a fragment thereof.

25. The nucleic acid of claim 24, wherein said fragment further comprises a nucleic acid encoding substantially the same nucleotide sequence as the variable region of said LM609 grafted light chain polypeptide (SEQ ID NO:31).

26. The nucleic acid of claim 24, wherein said fragment further comprises a nucleic acid encoding substantially the same nucleotide sequence as a CDR of said LM609 grafted light chain polypeptide.

27. A nucleic acid encoding a LM609 grafted antibody heavy chain polypeptide comprising a nucleotide sequence encoding substantially the same LM609 grafted heavy chain variable region amino acid sequence as that shown in Figure 1A (SEQ ID NO:2) or fragment thereof.

28. The nucleic acid of claim 27, wherein said fragment further comprises a nucleic acid encoding substantially the same heavy chain variable region amino acid sequence of said LM609 grafted heavy chain amino acid sequence (SEQ ID NO:2).

29. The nucleic acid of claim 27, wherein said fragment further comprises a nucleic acid encoding substantially the same heavy chain CDR amino acid sequence of said LM609 grafted heavy chain amino acid sequence.

30. A nucleic acid encoding a LM609 grafted antibody light chain polypeptide comprising a nucleotide sequence encoding substantially the same LM609 grafted light chain variable region amino acid sequence as that
5 shown in Figure 7 (SEQ ID NO:32) or fragment thereof.

31. The nucleic acid of claim 30, wherein said fragment further comprises a nucleic acid encoding substantially the same light chain variable region amino acid sequence of said Vitaxin light chain amino acid
10 sequence (SEQ ID NO:32).

32. The nucleic acid of claim 30, wherein said fragment further comprises a nucleic acid encoding substantially the same light chain CDR amino acid sequence of said Vitaxin light chain amino acid sequence.

33. A LM609 grafted heavy chain polypeptide comprising substantially the same variable region amino acid sequence as that shown in Figure 1A (SEQ ID NO:2) or functional fragment thereof.
15

34. The LM609 grafted heavy chain polypeptide of claim 33, wherein said functional fragment comprises a variable chain polypeptide or a CDR polypeptide.
20

35. A LM609 grafted light chain polypeptide comprising substantially the same variable region amino acid sequence as that shown in Figure 7 (SEQ ID NO:32) or
25 a functional fragment thereof.

36. The LM609 grafted light chain polypeptide of claim 35, wherein said functional fragment comprises a variable chain polypeptide or a CDR polypeptide.

260670-79097060

37. A nucleic acid encoding a heavy chain polypeptide for monoclonal antibody LM609 comprising substantially the same heavy chain variable region nucleotide sequence as that shown in Figure 2A (SEQ ID NO:5) or a fragment thereof.

38. The nucleic acid of claim 37, wherein said fragment further comprises a nucleic acid encoding substantially the same nucleotide sequence as the variable region of said heavy chain polypeptide (SEQ ID NO:5).

39. The nucleic acid of claim 37, wherein said fragment further comprises a nucleic acid encoding substantially the same nucleotide sequence as a CDR of said heavy chain polypeptide.

40. A nucleic acid encoding a light chain polypeptide for monoclonal antibody LM609 comprising substantially the same light chain variable region nucleotide sequence as that shown in Figure 2B (SEQ ID NO:7) or a fragment thereof.

41. The nucleic acid of claim 40, wherein said fragment further comprises a nucleic acid encoding substantially the same nucleotide sequence as the variable region of said light chain polypeptide (SEQ ID NO:7).

42. The nucleic acid of claim 40, wherein said fragment further comprises a nucleic acid encoding substantially the same nucleotide sequence as a CDR of said light chain polypeptide.

43. A nucleic acid encoding a heavy chain polypeptide for monoclonal antibody LM609 comprising a nucleotide sequence encoding substantially the same heavy chain variable domain amino acid sequence of monoclonal antibody LM609 as that shown in Figure 2A (SEQ ID NO:6) or fragment thereof.

44. The nucleic acid of claim 43, wherein said fragment further comprises a nucleic acid encoding substantially the same heavy chain variable region amino acid sequence of said monoclonal antibody LM609 (SEQ ID NO:6).

45. The nucleic acid of claim 43, wherein said fragment further comprises a nucleic acid encoding substantially the same heavy chain CDR amino acid sequence as said monoclonal antibody LM609.

46. A nucleic acid encoding a heavy chain polypeptide for monoclonal antibody LM609 comprising a nucleotide sequence encoding substantially the same light chain amino acid sequence of monoclonal antibody LM609 as that shown in Figure 2B (SEQ ID NO:8) or fragment thereof.

47. The nucleic acid of claim 46, wherein said fragment further comprises a nucleic acid encoding substantially the same light chain variable region amino acid sequence of said monoclonal antibody LM609 (SEQ ID NO:8).

48. The nucleic acid of claim 46, wherein said fragment further comprises a nucleic acid encoding substantially the same light chain CDR amino acid sequence as said monoclonal antibody LM609.

SECRET "T909T060

49. A method of inhibiting a function of $\alpha_v\beta_3$ comprising contacting $\alpha_v\beta_3$ with Vitaxin or a functional fragment thereof under conditions which allow binding of Vitaxin to $\alpha_v\beta_3$.

5 50. The method of claim 49, wherein said functional fragment is selected from the group consisting of Fv, Fab, F(ab)₂ and scFV.

51. The method of claim 49, wherein said function of $\alpha_v\beta_3$ is binding of $\alpha_v\beta_3$ to a ligand.

10 52. The method of claim 49, wherein said function of $\alpha_v\beta_3$ is integrin mediated signal transduction.

53. A method of treating an $\alpha_v\beta_3$ -mediated disease comprising administering an effective amount of Vitaxin or a functional fragment thereof under conditions
15 which allow binding to $\alpha_v\beta_3$.

54. The method of claim 53, wherein said functional fragment is selected from the group consisting of Fv, Fab, F(ab)₂ and scFV.

55. The method of claim 53, wherein said
20 $\alpha_v\beta_3$ -mediated disease is angiogenesis or restenosis.

56. An enhanced LM609 grafted antibody exhibiting selective binding affinity to $\alpha_v\beta_3$, or a functional fragment thereof, comprising at least one amino acid substitution in one or more CDRs of a LM609
25 grafted heavy chain variable region polypeptide or a LM609 grafted light chain variable region polypeptide, wherein the $\alpha_v\beta_3$ binding affinity of said enhanced LM609 grafted antibody is maintained.

0304001-0399

57. The enhanced LM609 grafted antibody of claim 56, wherein said $\alpha_v\beta_3$ binding affinity of said LM609 grafted antibody is enhanced.

58. The enhanced LM609 grafted antibody of
5 claim 56, wherein said functional fragment is selected from the group consisting of Fv, Fab, F(ab)₂ and scFV.

59. The enhanced LM609 grafted antibody of claim 56, wherein said CDR having at least one amino acid substitution is selected from the group consisting of
10 V_H CDR1, V_H CDR2, V_H CDR3, V_L CDR1, V_L CDR2 and V_L CDR3.

60. The enhanced LM609 grafted antibody of claim 59, wherein said V_H CDR1 is selected from the group consisting of the CDRs referenced as SEQ ID NO:48, SEQ ID NO:50 and SEQ ID NO:52.

61. The enhanced LM609 grafted antibody of
15 claim 59, wherein said V_H CDR2 is selected from the group consisting of the CDRs referenced as SEQ ID NO:54, SEQ ID NO:56 and SEQ ID NO:58.

62. The enhanced LM609 grafted antibody of
20 claim 59, wherein said V_H CDR3 is selected from the group consisting of the CDRs referenced as SEQ ID NO:60, SEQ ID NO:62, SEQ ID NO:64, SEQ ID NO:66, SEQ ID NO:68, SEQ ID NO:70, SEQ ID NO:72, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:94, SEQ ID NO:96; SEQ ID
25 NO:98 and SEQ ID NO:100.

63. The enhanced LM609 grafted antibody of claim 59, wherein said V_L CDR1 is the CDR referenced as SEQ ID NO:82.

SEQUENCE LISTING

64. The enhanced LM609 grafted antibody of claim 59, wherein said V_L CDR2 is the CDR referenced as SEQ ID NO:84.

65. The enhanced LM609 grafted antibody of claim 59, wherein said V_L CDR3 is selected from the group consisting of the CDRs referenced as SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:90 and SEQ ID NO:92.

66. The enhanced LM609 grafted antibody of claim 56, wherein said enhanced LM609 grafted antibody comprises at least one amino acid substitution in two or more CDRs of a LM609 grafted heavy chain variable region polypeptide or a LM609 grafted light chain variable region polypeptide.

67. The enhanced LM609 grafted antibody of claims 66, wherein said functional fragment is selected from the group consisting of Fv, Fab, $F(ab)_2$ and scFV.

68. The enhanced LM609 grafted antibody of claim 66, wherein said CDR having at least one amino acid substitution is selected from the group consisting of V_H CDR1, V_H CDR2, V_H CDR3, V_L CDR1, V_L CDR2 and V_L CDR3.

69. The enhanced LM609 grafted antibody of claim 68, wherein said enhanced LM609 grafted antibody comprises the combination of CDRs selected from the group consisting of:

- 5 the V_L CDR1 referenced as SEQ ID NO:82 and the V_H CDR3 referenced as SEQ ID NO:68;
- the V_L CDR1 referenced as SEQ ID NO:82, the V_H CDR2 referenced as SEQ ID NO:56 and the V_H CDR3 referenced as SEQ ID NO:68;
- 10 the V_L CDR1 referenced as SEQ ID NO:82, the V_H CDR2 referenced as SEQ ID NO:56 and the V_H CDR3 referenced as SEQ ID NO:72;
- the V_L CDR1 referenced as SEQ ID NO:82, the V_H CDR2 referenced as SEQ ID NO:56 and the V_H CDR3 referenced as SEQ ID NO:70;
- 15 the V_L CDR1 referenced as SEQ ID NO:82 and the V_H CDR3 referenced as SEQ ID NO:72;
- the V_L CDR3 referenced as SEQ ID NO:86, the V_H CDR2 referenced as SEQ ID NO:56 and the V_H CDR3 referenced as SEQ ID NO:68;
- 20 the V_L CDR3 referenced as SEQ ID NO:90 and V_H CDR3 referenced as SEQ ID NO:68; and
- the V_L CDR3 referenced as SEQ ID NO:90, the V_H CDR2 referenced as SEQ ID NO:56 and V_H CDR3 referenced as SEQ ID NO:68.
- 25

70. The enhanced LM609 grafted antibody of claim 66, wherein at least one of said CDRs has two or more amino acid substitutions.

71. The enhanced LM609 grafted antibody of claims 70, wherein said functional fragment is selected from the group consisting of Fv, Fab, F(ab)₂ and scFV.

30

860670 "T909T060

72. The enhanced LM609 grafted antibody of claim 70, wherein said CDR having at least one amino acid substitution is selected from the group consisting of V_H CDR1, V_H CDR2, V_H CDR3, V_L CDR1, V_L CDR2 and V_L CDR3.

5 73. The enhanced LM609 grafted antibody of
claim 72, wherein said enhanced LM609 grafted antibody
comprises the combination of CDRs selected from the group
consisting of:

the V_L CDR1 referenced as SEQ ID NO:82, the V_H
10 CDR2 referenced as SEQ ID NO:56 and the V_H CDR3 referenced
as SEQ ID NO:94;

the V_L CDR3 referenced as SEQ ID NO:90, the V_H CDR2 referenced as SEQ ID NO:56 and the V_H CDR3 referenced as SEQ ID NO:94;

15 the V_L CDR3 referenced as SEQ ID NO:90, the V_H
CDR2 referenced as SEQ ID NO:56 and the V_H CDR3 referenced
as SEQ ID NO:96;

the V_L CDR3 referenced as SEQ ID NO:90 and the V_H CDR3 referenced as SEQ ID NO:94;

20 the V_L CDR3 referenced as SEQ ID NO:90 and the
V_H CDR3 referenced as SEQ ID NO:98; and

the V_L CDR3 referenced as SEQ ID NO:90, the V_H CDR2 referenced as SEQ ID NO:56 and the V_H CDR3 referenced as SEQ ID NO:100.

25 74. A high affinity LM609 grafted antibody exhibiting selective binding affinity to $\alpha_v\beta_3$, or a functional fragment thereof, comprising at least one amino acid substitution in one or more CDRs of a LM609 grafted heavy chain variable region polypeptide or a
30 LM609 grafted light chain variable region polypeptide, wherein the $\alpha_v\beta_3$ binding affinity of said high affinity LM609 grafted antibody is enhanced.

75. The high affinity LM609 grafted antibody of claim 74, wherein said functional fragment is selected from the group consisting of Fv, Fab, F(ab)₂ and scFV.

76. The high affinity LM609 grafted antibody
5 of claim 74, wherein said CDR having at least one amino acid substitution is selected from the group consisting of V_H CDR1, V_H CDR2, V_H CDR3, V_L CDR1, V_L CDR2 and V_L CDR3.

77. The high affinity LM609 grafted antibody
10 of claim 76, wherein said high affinity LM609 grafted antibody comprises the combination of CDRs selected from the group consisting of:

the V_L CDR1 referenced as SEQ ID NO:82 and the V_H CDR3 referenced as SEQ ID NO:68;

the V_L CDR1 referenced as SEQ ID NO:82, the V_H
15 CDR2 referenced as SEQ ID NO:56 and the V_H CDR3 referenced as SEQ ID NO:68;

the V_L CDR1 referenced as SEQ ID NO:82, the V_H
CDR2 referenced as SEQ ID NO:56 and the V_H CDR3 referenced as SEQ ID NO:72;

20 the V_L CDR1 referenced as SEQ ID NO:82, the V_H CDR2 referenced as SEQ ID NO:56 and the V_H CDR3 referenced as SEQ ID NO:70;

the V_L CDR1 referenced as SEQ ID NO:82 and the V_H CDR3 referenced as SEQ ID NO:72;

25 the V_L CDR3 referenced as SEQ ID NO:86, the V_H CDR2 referenced as SEQ ID NO:56 and the V_H CDR3 referenced as SEQ ID NO:68;

the V_L CDR3 referenced as SEQ ID NO:90, the V_H CDR2 referenced as SEQ ID NO:56 and the V_H CDR3 referenced
30 as SEQ ID NO:94;

the V_L CDR3 referenced as SEQ ID NO:90 and V_H CDR3 referenced as SEQ ID NO:68;

the V_L CDR3 referenced as SEQ ID NO:90, the V_H CDR2 referenced as SEQ ID NO:56 and V_H CDR3 referenced as SEQ ID NO:68;

the V_L CDR1 referenced as SEQ ID NO:82, the V_H CDR2 referenced as SEQ ID NO:56 and the V_H CDR3 referenced as SEQ ID NO:94;

the V_L CDR3 referenced as SEQ ID NO:90, the V_H CDR2 referenced as SEQ ID NO:56 and the V_H CDR3 referenced as SEQ ID NO:96;

the V_L CDR3 referenced as SEQ ID NO:90 and the V_H CDR3 referenced as SEQ ID NO:94;

the V_L CDR3 referenced as SEQ ID NO:90 and the V_H CDR3 referenced as SEQ ID NO:98; and

the V_L CDR3 referenced as SEQ ID NO:90, the V_H CDR2 referenced as SEQ ID NO:56 and the V_H CDR3 referenced as SEQ ID NO:100.

78. A nucleic acid encoding an enhanced LM609 grafted antibody, or a functional fragment thereof, exhibiting selective binding affinity to $\alpha_v\beta_3$ comprising at least one amino acid substitution in one or more CDRs of a LM609 grafted heavy chain variable region polypeptide or a LM609 grafted light chain variable region polypeptide, wherein the $\alpha_v\beta_3$ binding affinity of said enhanced LM609 grafted antibody is maintained or enhanced.

79. A nucleic acid encoding a high affinity LM609 grafted antibody, or a functional fragment thereof, exhibiting selective binding affinity to $\alpha_v\beta_3$ comprising at least one amino acid substitution in one or more CDRs of a LM609 grafted heavy chain variable region polypeptide or a LM609 grafted light chain variable region polypeptide, wherein the $\alpha_v\beta_3$ binding affinity of said high affinity LM609 grafted antibody is enhanced.

ABSTRACT OF THE INVENTION

The invention provides a Vitaxin antibody and a LM609 grafted antibody exhibiting selective binding affinity to $\alpha_v\beta_3$. The Vitaxin antibody consists of at least one Vitaxin heavy chain polypeptide and at least one Vitaxin light chain polypeptide or functional fragments thereof. Also provided are the Vitaxin heavy and light chain polypeptides and functional fragments. The LM609 grafted antibody consists of at least one CDR grafted heavy chain polypeptide and at least one CDR grafted light chain polypeptide or functional fragment thereof. The invention additionally provides a high affinity LM609 grafted antibody comprising one or more CDRs having at least one amino acid substitution, where the $\alpha_v\beta_3$ binding activity of the high affinity LM609 grafted antibody is enhanced. Nucleic acids encoding Vitaxin and LM609 grafted heavy and light chains as well as nucleic acids encoding the parental non-human antibody LM609 are additionally provided. Functional fragments of such encoding nucleic acids are similarly provided. The invention also provides a method of inhibiting a function of $\alpha_v\beta_3$. The method consists of contacting $\alpha_v\beta_3$ with Vitaxin or a LM609 grafted antibody or functional fragments thereof under conditions which allow binding to $\alpha_v\beta_3$. Finally, the invention provides for a method of treating an $\alpha_v\beta_3$ -mediated disease. The method consists of administering an effective amount of Vitaxin or a LM609 grafted antibody or functional fragment thereof under conditions which allow binding to $\alpha_v\beta_3$.

[illegible]

[illegible]

3/13

Figure 2a

GAA	GTG	CAG	CTG	GTG	GAG	TCT	GGG	GGA	GGC	TTA	GTG	AAG	CCT	GGA	AGG	48
Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Lys	Pro	Gly	Arg	
1				5					10					15		
TCC	CTG	AGA	CTC	TCC	TGT	GCA	GCC	TCT	GGA	TTC	GCT	TTC	AGT	AGC	TAT	96
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Ala	Phe	Ser	Ser	Tyr	
			20					25					30			
GAC	ATG	TCT	TGG	GTT	CGC	CAG	ATT	CCG	GAG	AAG	AGG	CTG	GAG	TGG	GTC	144
Asp	Met	Ser	Trp	Val	Arg	Gln	Ile	Pro	Glu	Lys	Arg	Leu	Glu	Trp	Val	
		35					40					45				
GCA	AAA	GTT	AGT	AGT	GGT	GGT	GGT	AGC	ACC	TAC	TAT	TTA	GAC	ACT	GTG	192
Ala	Lys	Val	Ser	Ser	Gly	Gly	Gly	Ser	Thr	Tyr	Tyr	Leu	Asp	Thr	Val	
	50					55					60					
CAG	GGC	CGA	TTC	ACC	ATC	TCC	AGA	GAC	AAT	GCC	AAG	AAC	ACC	CTA	TAC	240
Gln	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Asn	Thr	Leu	Tyr	
65					70					75					80	
CTG	CAA	ATG	AGC	AGT	CTG	AAC	TCT	GAG	GAC	ACA	GCC	ATG	TAT	TAC	TGT	288
Leu	Gln	Met	Ser	Ser	Leu	Asn	Ser	Glu	Asp	Thr	Ala	Met	Tyr	Tyr	Cys	
				85					90					95		
GCA	AGA	CAT	AAC	TAC	GGC	AGT	TTT	GCT	TAC	TGG	GGC	CAA	GGG	ACT	CTG	336
Ala	Arg	His	Asn	Tyr	Gly	Ser	Phe	Ala	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	
			100					105					110			
GTC	ACT	GTC	TCT	GCA												351
Val	Thr	Val	Ser	Ala												
			115													

05016061-013098

4/13

Figure 2b

GAT	ATT	GTG	CTA	ACT	CAG	TCT	CCA	GCC	ACC	CTG	TCT	GTG	ACA	CCA	GGA	48
Asp	Ile	Val	Leu	Thr	Gln	Ser	Pro	Ala	Thr	Leu	Ser	Val	Thr	Pro	Gly	
1				5					10					15		
GAT	AGC	GTC	AGT	CTT	TCC	TGC	CAG	GCC	AGC	CAA	AGT	ATT	AGC	AAC	CAC	96
Asp	Ser	Val	Ser	Leu	Ser	Cys	Gln	Ala	Ser	Gln	Ser	Ile	Ser	Asn	His	
			20					25					30			
CTA	CAC	TGG	TAT	CAA	CAA	AAA	TCA	CAT	GAG	TCT	CCA	AGG	CTT	CTC	ATC	144
Leu	His	Trp	Tyr	Gln	Gln	Lys	Ser	His	Glu	Ser	Pro	Arg	Leu	Leu	Ile	
		35				40					45					
AAG	TAT	CGT	TCC	CAG	TCC	ATC	TCT	GGG	ATC	CCC	TCC	AGG	TTC	AGT	GGC	192
Lys	Tyr	Arg	Ser	Gln	Ser	Ile	Ser	Gly	Ile	Pro	Ser	Arg	Phe	Ser	Gly	
	50					55				60						
AGT	GGA	TCA	GGG	ACA	GAT	TTC	GCT	CTC	AGT	ATC	AAC	AGT	GTG	GAG	ACT	240
Ser	Gly	Ser	Gly	Thr	Asp	Phe	Ala	Leu	Ser	Ile	Asn	Ser	Val	Glu	Thr	
65					70					75				80		
GAA	GAT	TTT	GGA	ATG	TAT	TTC	TGT	CAA	CAG	AGT	GGC	AGC	TGG	CCT	CAC	288
Glu	Asp	Phe	Gly	Met	Tyr	Phe	Cys	Gln	Gln	Ser	Gly	Ser	Trp	Pro	His	
				85				90						95		
ACG	TTC	GGA	GGG	GGG	ACC	AAG	CTG	GAA	ATT	AAG						321
Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys						
			100					105								

09016061 013098

5/13

LM609 Competition Assay




Figure 3

6/13

Figure 4A




Figure 4B





Figure 4C



7/13

Figure 5A





Figure 5B



8/13

Figure 6A




Figure 6B




Figure 6C




Figure 7

GAG Glu 1	ATT Ile	GTG Val	CTA Leu	ACT Thr 5	CAG Gln	TCT Ser	CCA Pro	GCC Ala	ACC Thr 10	CTG Leu	TCT Ser	CTC Leu	AGC Ser	CCA Pro 15	GGA Gly	48
GAA Glu	AGG Arg	GCG Ala	ACT Thr 20	CTT Leu	TCC Ser	TGC Cys	CAG Gln	GCC Ala 25	AGC Ser	CAA Gln	AGT Ser	ATT Ile	AGC Ser 30	AAC Asn	CAC His	96
CTA Leu	CAC His	TGG Trp 35	TAT Tyr	CAA Gln	CAA Gln	AGG Arg	CCT Pro 40	GGT Gly	CAA Gln	GCC Ala	CCA Pro	AGG Arg 45	CTT Leu	CTC Leu	ATC Ile	144
CGT/ATG Arg/Met	TAT Tyr 50	CGT Arg	TCC Ser	CAG Gln	TCC Ser	ATC Ile 55	TCT Ser	GGG Gly	ATC Ile	CCC Pro	GCC Ala 60	AGG Arg	TTC Phe	AGT Ser	GGC Gly	192
AGT Ser 65	GGA Gly	TCA Ser	GGG Gly	ACA Thr	GAT Asp 70	TTC Phe	ACC Thr	CTC Leu	ACT Thr 75	ATC Ile	TCC Ser	AGT Ser	CTG Leu	GAG Glu	CCT Pro 80	240
GAA Glu	GAT Asp	TTT Phe	GCA Ala	GTC Val 85	TAT Tyr	TAC Tyr	TGT Cys	CAA Gln 90	CAG Gln	AGT Ser	GGC Gly	AGC Ser	TGG Trp 95	CCT Pro	CAC His	288
ACG Thr	TTC Phe	GGA Gly	GGG Gly 100	GGG Gly	ACC Thr	AAG Lys	GTG Val	GAA Glu 105	ATT Ile	AAG Lys						321

10/13




FIGURE 8

11/13




FIGURE 9

SECRET 79097060

12/13




FIGURE 10

13/13





FIGURE 11



Application deficiencies found during scanning:

1. Application papers are not suitable for scanning and are not in compliance with 37 CFR 1.52 because:
- ☐ All sheets must be the same size and either A4 (21 cm x 29.7 cm) or 8-1/2" x 11". Pages _____ do not meet these requirements.
 - ☐ Papers are not flexible, strong, smooth, non-shiny, durable, and white.
 - ☐ Papers are not typewritten or mechanically printed in permanent ink on one side.
 - ☐ Papers contain improper margins. Each sheet must have a left margin of at least 2.5 cm (1") and top, bottom and right margins of at least 2.0 cm (3/4").
 - ☐ Papers contain hand lettering.
2. Drawings are not in compliance and were not scanned because:
- ☐ The drawings or copy of drawings are not suitable for electronic reproduction.
 - ☐ All drawings sheets are not the same size. Pages must be either A4 (21 cm x 29.7 cm) or 8-1/2" x 11".
 - ☐ Each sheet must include a top and left margin of at least 2.5 cm (1"), a right margin of at least 1.5 cm (9/16") and a bottom margin of at least 1.0 cm (3/8").
3. Page(s) _____ are not of sufficient clarity, contrast and quality for electronic reproduction.
4. Page(s) _____ are missing.
5. OTHER: No Declaration

05016061-0130980